

Increased sulfate uptake by *E. coli* overexpressing the SLC26-related SulP protein Rv1739c from *Mycobacterium tuberculosis*

Alexander S. Zolotarev^a, Meera Unnikrishnan^b, Boris E. Shmukler^a, Jeffrey S. Clark^a,
David H. Vidorpe^a, Nikolaus Grigorieff^{c,d}, Eric J. Rubin^{b,e,f}, Seth L. Alper^{a,f,*}

^a Molecular and Vascular Medicine and Renal Divisions, Beth Israel Deaconess Medical Center, USA

^b Department of Immunology and Infectious Diseases, Harvard School of Public Health, USA

^c Howard Hughes Medical Institute, Brandeis University, Waltham, MA, USA

^d Department of Biochemistry, Brandeis University, Waltham, MA, USA

^e Departments of Microbiology and Molecular Genetics, Harvard Medical School, Boston, MA, USA

^f Department Medicine, Harvard Medical School, Boston, MA, USA

Received 15 November 2007; received in revised form 13 December 2007; accepted 14 December 2007

Available online 23 December 2007

Abstract

Growth and virulence of mycobacteria requires sulfur uptake. The *Mycobacterium tuberculosis* genome contains, in addition to the ABC sulfate permease *cysTWA*, three SLC26-related SulP genes of unknown function. We report that induction of Rv1739c expression in *E. coli* increased bacterial uptake of sulfate, but not Cl^- , formate, or oxalate. Uptake was time-dependent, maximal at pH 6.0, and exhibited a $K_{1/2}$ for sulfate of 4.0 μM . Na^+ -independent sulfate uptake was not reduced by bicarbonate, nitrate, or phosphate, but was inhibited by sulfite, selenate, thiosulfate, N-ethylmaleimide and carbonyl cyanide 3-chloro-phenylhydrazone. Sulfate uptake was also increased by overexpression of the Rv1739c transmembrane domain, but not of the cytoplasmic C-terminal STAS domain. Mutation to serine of the three cysteine residues of Rv1739c did not affect magnitude, pH-dependence, or pharmacology of sulfate uptake. Expression of Rv1739c in a *M. bovis* BCG strain lacking the ABC sulfate permease subunit CysA could not complement sulfate auxotrophy. Moreover, inducible expression of Rv1739c in an *E. coli* strain lacking CysA did not increase sulfate uptake by intact cells. Our data show that facilitation of bacterial sulfate uptake by Rv1739c requires CysA and its associated sulfate permease activity, and suggest that Rv1739c may be a CysTWA-dependent sulfate transporter.

© 2007 Elsevier Inc. All rights reserved.

Keywords: *Mycobacterium tuberculosis*; *E. coli*; *M. bovis*; Sulfate transport; SLC26A6; SulP

1. Introduction

Eukaryotic bicarbonate/anion transporters are encoded by the SLC4 and SLC26 gene superfamilies. Inherited human diseases are associated with mutations in multiple members of both superfamilies (Mount and Romero, 2004; Romero et al., 2004; Alper, 2006; Kere, 2006). However, the transmembrane domain structure of neither family is currently known. SLC26-related SulP genes are widely distributed among bacterial species (Felce and Saier, 2004; Price et al., 2004), in addition to the widespread

expression of SLC26 genes among eukaryotic organisms (Mount and Romero, 2004). Vertebrate SLC26 polypeptides transport a wide range of anions, including Cl^- , HCO_3^- , iodide, formate, oxalate, and sulfate, and have been implicated in teleost kidney sulfate excretion and osmoregulation (Renfro et al., 1999; Nakada et al., 2005; Katoh et al., 2006). SulP sulfate transporters of plants and yeast have been characterized functionally, and evidence of SulP-mediated HCO_3^- transport has been demonstrated in *Synechococcus* species (Price et al., 2004), but little other functional data on bacterial SulP proteins is available. Bacterial proteins have been important for catalyzing progress in the structure determination of polytopic membrane proteins of higher organisms. Functional data on prokaryotic SulP proteins is important for purification of native protein for crystallization and other structural studies.

* Corresponding author. Molecular and Vascular Medicine and Renal Divisions, E/RW763 Beth Israel Deaconess Medical Center, 330 Brookline Avenue, Boston, MA 02215, USA.

E-mail address: salper@bidmc.harvard.edu (S.L. Alper).




Fig. 1. Overexpression of Rv1739c increases sulfate uptake. A. Schematic of Rv1739c polypeptide from *M. tuberculosis*. The hypothetical disposition of the transmembrane spans is based on hydropathy analysis described by (Felce and Saier, 2004). B. Left: Anti-6His immunoblot of clarified SDS lysate of uninduced (lane 1) and IPTG-induced Tuner(DE3)pLacI cells (lane 2). Right: Coomassie Blue R250-stained gel of Rv1739c eluted from Ni-NTA column by 250 mM imidazole. C. ³⁵S-sulfate uptake in uninduced Tuner cells or cells induced to express either *M. tuberculosis* Rv1739c (left bars) or *E. coli* ychM (right bars). Number of measurements is indicated within parentheses. *, $p < 0.05$. D. Anti-6His immunoblot of clarified SDS lysate of uninduced Tuner cells (lane 1) or Tuner cells induced to express ychM polypeptide (lane 2).

As is true for eukaryotes, sulfur is a key element in bacterial metabolism. Rapidly growing numbers of anaerobic, sulfate-reducing chemolithoautotrophic species have been identified in samples from deep sea hydrothermal vents, both as free-living bacteria and as thiotrophic endo- and exosymbionts of deep vent marine invertebrates (Rosenberg et al., 2006). Genes involved in sulfur metabolism have been implicated as virulence determinants in mammalian pathogens. In mycobacteria, mycothiols have been reported to contribute to anti-oxidant defense (Fan et al., 2007). Sulfolipids present in the mycobacterial outer envelope act as both positive and negative regulators of *M. tuberculosis* pathogenicity (Schelle and Bertozzi, 2006). Overexpression screening of several bacterial SulP genes revealed inducible, stable accumulation of *M. tuberculosis* Rv1739c and *E. coli* ychM. In view of the importance of the SulP gene family in plant (Yoshimoto et al., 2007) and yeast (Shibagaki and Grossman, 2006), and fish biology (Renfro et al., 1999; Nakada et al., 2005; Katoh et al., 2006), the significance of the SLC26 gene superfamily in mammalian biology (Mount and Romero, 2004; Markovich and Aronson, 2007), and the proposed importance of sulfate in *M. tuberculosis* pathogenicity (Schelle and Bertozzi, 2006; Fan et al., 2007), we investigated further the function of Rv1739c (Fig. 1A) as a candidate sulfate transporter of mycobacteria.

2. Materials and methods

2.1. Molecular cloning

M. tuberculosis H37Rv genomic cosmid MTCY4C12 containing the Rv1739c gene (NC_000962) was provided by K. Eiglmeier (Inst. Pasteur, Paris). Rv1739c DNA encoding the predicted 560 aa Rv1739c SulP polypeptide (CAB03701) was PCR-amplified from the cosmid with addition of a C-terminal 6His tag. *E. coli* K-12 MG1655 BamHI genomic clone DD267 in phage λ Charon was provided by the G. Plunkett and F. Blattner (*E. coli* Genome Project, Univ. Wisconsin). ychM DNA encoding the (predicted) 550 aa¹ ychM SulP polypeptide

¹ The ychM open reading frame of 550 aa has been annotated as initiating from an AUG codon in *E. coli* O157:H7 (BAB35134) and in *E. coli* APEC 01 (YP_852337). The ychM open reading frame has also been annotated as 559 aa, initiating from an inframe upstream GUG codon in *E. coli* W3110 (AP_001831) and K12 (NP_415724). The overlapping portions of the two amino acid sequences are identical. Yet another possible initiation site is the GUG initiation codon giving rise to a predicted ychM ORF of 551 aa, and the corresponding initiation site has been annotated in the closely related *S. typhimurium* ychM (AAL20696; 553 aa, 93% aa identity with *E. coli* ychM). None of the possible translational initiation sites has been validated experimentally. We amplified the 550 aa ORF of ychM, since the overexpression of the 559 aa ORF (JW5189) was reported by the ASKA collection curators to be toxic (<http://ecoli.naist.jp/GB6/info.jsp?id=JW5189>), and was among the polypeptides that could not be purified as a bait protein in a global protein-protein interaction screen (Arifuzzaman et al., 2006).

(BAB35134) was PCR-amplified from the phage DNA, also with addition of a C-terminal 6His tag. Similar approaches were used to generate DNAs encoding the isolated Rv1739c transmembrane domain (aa 1–436, in which aa 411–436 are predicted to reside in the juxtamembrane cytosol) and the isolated Rv1739c C-terminal cytoplasmic region encompassing its STAS domain (aa 437–560). PCR products were subcloned into pETBlue-1 (Perfectly Blunt Cloning Kit, Novagen) and sequenced on both strands. pETBlue-1-SulP constructs were transformed into Tuner DE3pLacI cells (Novagen) for inducible expression. Oligonucleotide primers used are listed in Supplemental Table 1.

The Cys-less Rv1739c was constructed with QuickChange Multi Site-Directed Mutagenesis Kit (Stratagene). Directed mutagenesis of single codons was performed with the QuickChange II Site-Directed Mutagenesis Kit (Stratagene). PCR was used to introduce the Kozak consensus sequence CACTCCCG-CAGG from mouse Ae2b1 (Lecanda et al., 2000) immediately 5' to the Rv1739c initiator methionine, and the modified DNA was subcloned into the *Xenopus* oocyte expression vector pXT7. This Kozak sequence is active in *Xenopus* oocytes (Kurschat et al., 2006).

Wildtype *E. coli* strain BW25113 (obtained from Yale *E. coli* Genetic Stock Center) and its *cysA* deletion mutant JD21556 (obtained from National Bioresource Project of the Institute of Genetics, Japan; <http://www.shigen.nig.ac.jp/ecoli/strain/transposoneDetailAction.do?strainId=1187>) were both grown without antibiotics. Rv1739c-6His was subcloned into pBAD33 (Guzman et al., 1995) for expression in JD21556. Rv1739c-6His was also subcloned into pMV261.hyg for GroEL promoter-driven constitutive expression in the wildtype *M. bovis* BCG strain, grown in 7H9T liquid medium (Wooff et al., 2002). The BCG *cysA* deletion strain EWP44 (the kind gift of Esen Wooff, Veterinary Laboratories Agency, Addlestone, Surrey, UK), was grown in 7H9T liquid medium supplemented with kanamycin and methionine (Wooff et al., 2002). *M. bovis* expression plasmid pSM-101EW encoding *M. tuberculosis* CysA (Wooff et al., 2002) was also the gift of Esen Wooff. Bacterial strains are summarized in Supplemental Table 2.

All subclones generated by PCR were verified by DNA sequencing of both strands, and by immunoblot confirmation of IPTG-induced biosynthesis of 6His-tagged polypeptide of the appropriate M_r using anti-6His monoclonal antibody ("His-Tag", Novagen).

2.2. Isotopic anion influx measurement

An 8 h culture of Tuner DE3pLacI cells transformed with pET-Blue-1-Rv1739c-6His or with pET-Blue-1-yehM-6His picked from a single colony was grown in LB supplemented with 1% glucose, 50 μ g/ml carbenicillin, and 30 μ g/ml chloramphenicol. 50 ml fresh LB medium with or without 0.5 mM isopropylthiogalactoside (IPTG) was inoculated with an aliquot of liquid culture and grown overnight at 20–22 °C with shaking at 300 rpm. Cultures were harvested by centrifugation upon reaching O.D. 0.3–0.5.

35 S-sulfate uptake into bacterial suspensions was measured as described (Lindblow-Kull et al., 1985), with modifications. *E. coli* suspensions in pre-weighed 50 ml Falcon tubes were pelleted and washed once in 10–15 ml "uptake medium" containing (in mM) 62 K phosphate, 47 NH_4Cl , 1.7 Na citrate, 0.6 MgCl_2 , and 11 glucose, pH 7.0. Washed pellets were resuspended at 25 mg/ml in uptake medium supplemented with 0.5 mM Na_2SO_4 , incubated 30 min at room temp, pelleted, and resuspended in identical volumes of uptake medium containing 0.1 mM sulfate or the indicated concentrations at indicated pH, with additional agents as indicated. Uptake was initiated by addition of 1–2 $\mu\text{Ci/ml}$ ^{35}S -sulfate (ICN) with vortexing. 0.1 ml samples were removed at 0 and 5 min, or at the times indicated, rapidly filtered through Millipore GSTF02500 filters mounted on a manifold (Hoeffer FH224V). Filters were immediately washed twice with 2.5 ml sulfate-free uptake medium and, after drying, ^{35}S was measured by scintillation counting. All uptake experiments were performed at room temperature. All uptake data presented in individual figure panels represents experiments in which uninduced and induced bacteria were prepared in parallel and assayed the same day.

2.3. *Xenopus* oocyte expression and assays

crRNA encoding the Rv1739c open reading frame preceded by the Kozak initiation sequence from mouse Kcc1/Slc12a1 (Casula et al., 2001) was subcloned into the *Xenopus* oocyte transcription vector, pXT7.

^{35}S -sulfate influx and two-electrode voltage clamp currents were assayed as previously described (Chernova et al., 2005; Clark et al., in press).

2.4. Assay for complementation of *M. bovis* BCG auxotrophy

Competent cells of the *cysA* deletion strain EWP44 (Wooff et al., 2002) were electroporated with pMV261.hyg Rv1739c or with the empty plasmid, and plated on 7H10 agar (Wooff et al., 2002) with kanamycin and hygromycin in the presence or absence of methionine. EWP44 was also electroporated with the CysA-expressing plasmid pSM101EW and plated on 7H10 agar with kanamycin, in the presence or absence of methionine. Growth was evaluated ~3 weeks later.

3. Results

3.1. Induced overexpression of *M. tuberculosis* Rv1739c, but not of *E. coli* yehM, increases sulfate uptake by *E. coli*

IPTG-induction of the *M. tuberculosis* SulP gene Rv1739c in *E. coli* increased abundance of the 560 aa Rv1739c polypeptide (Fig. 1A) as detected by 6His-immunoblot of lysate and by Coomassie stain of Ni-NTA-enriched protein (Fig. 1B). The Rv1739c polypeptide migrated on SDS-PAGE as expected for a monomer. Increased expression of Rv1739c was accompanied by increased sulfate uptake into intact bacteria (Fig. 1C). Increased sulfate uptake did not accompany increased expression of yehM at similar abundance (Fig. 1C and D). As




Fig. 2. A. Growth phase-dependence of ^{35}S -sulfate uptake by native Tuner cells untreated with IPTG. B. Time-dependence of ^{35}S -sulfate uptake by uninduced Tuner cells (diamonds) and by cells induced to express Rv1739c (squares). C. ^{35}S -sulfate uptake into uninduced cells and cells induced to express Rv1739c. D. Sulfate uptake into cells harboring (Rv1739c) or lacking the Rv1739c expression plasmid (Tuner), after treatment without (uninduced) or with IPTG (induced). Number of measurements is indicated within parentheses. *, $p < 0.05$.

evident in the immunoblots, the uninduced T7 promoter of the pETBlue-1 vector exhibited leaky expression of both Rv1739c and ychM.

3.2. Tuner DE3pLacI cells exhibit strong density-dependent induction of native sulfate uptake

While optimizing the sulfate uptake assay, we noted that Tuner DE3pLacI cells exhibit strong upregulation of sulfate uptake in high density liquid culture (Fig. 2A). Rv1739c-expressing bacteria harvested from saturation culture did not

exhibit an increase of sulfate uptake over this high background of saturation-cultured cells. However, cultures at early-mid log phase (O.D. 0.3–0.5) reliably exhibited stimulation of sulfate uptake associated with induction of Rv1739c expression. This Rv1739c-associated sulfate uptake was time-dependent, roughly linear over 30 min (Fig. 2B), highly reproducible (Fig. 2C), and was not evident in untransformed Tuner cells exposed to IPTG (Fig. 2D). The magnitude of sulfate uptake was not altered by omission of the 30 min loading in 0.5 mM Na sulfate prior to the influx assay or by a 30 min pre-incubation period in the absence of sulfate ($n=4$, not shown).




Fig. 3. A. pH-dependence of ^{35}S -sulfate uptake into cells induced to express Rv1739c. Number of measurements is indicated within parentheses. *, $p < 0.05$. B. Sulfate concentration dependence of ^{35}S -sulfate uptake into cells expressing Rv1739c ($n=6$), fit with a hyperbolic equation (SigmaPlot).




Fig. 4. Tests of inhibition of ³⁵S-sulfate uptake into cells expressing Rv1739c by (A) 1 mM NEM, (B) 50 μM CCCP, (C) 5 mM Na sulfite, (D) 5 mM Na selenate, (E) 5 mM Na molybdate or Na thiosulfate, (F) 1 mM DIDS or 100 μM niflumate. ³⁵S-sulfate concentration was 100 μM in the presence of sulfite, molybdate, or thiosulfate, and 50 μM in the presence of the other antagonists. Number of measurements is within parentheses.

3.3. Rv1739c-associated sulfate uptake is maximal at pH 6 and is of high apparent affinity

Rv1739c-associated sulfate uptake was inhibited at pH 8.0 and maximal at pH 6.0 (Fig. 3A), but nearly abolished by further acidification to pH 5.0. Sulfate uptake in uninduced Tuner cells transformed showed qualitatively similar changes at low background levels. Rv1739c-associated sulfate influx exhibited a $K_{1/2}$ for extracellular sulfate of $4.04 \mu\text{M}$ (Fig. 3B). Tuner cells induced to express Rv1739c did not exhibit increased uptake of $^{36}\text{Cl}^-$, ^{14}C -formate, ^{14}C -oxalate at pH 6.0 or pH 7.0 (not shown). Thus, Rv1739c-associated sulfate uptake was of high affinity, maximal at pH 6, and sulfate-specific.

As rare examples of bacterial transporters have been reported to function when expressed in *Xenopus* oocytes (Quick and Wright, 2002), Rv1739c was subcloned into the oocyte transcription vector pXT7 behind the Kozak sequence from mouse *Kcc1/Slc12a1* (Casula et al., 2001), and functionally tested in *Xenopus* oocytes for ^{35}S -sulfate influx and efflux. The 30 min sulfate uptake into individual oocytes from ND-96 medium at pH 6.0 was slightly increased over background ($0.23 \pm 0.01 \text{ pmol min}^{-1}$, $n=30$) by Rv1739c expression in oocytes injected with 10 ng RNA ($0.31 \pm 0.02 \text{ pmol min}^{-1}$, $n=48$, $p < 0.005$), but not at pH 7.4 ($n=30$). However, interpretation of this cRNA-associated stimulation was confounded by lack of increased sulfate uptake into oocytes injected with 25 ng Rv1739c cRNA ($n=30$). Moreover, oocytes injected with 10 ng cRNA exhibited no sulfate-induced current at extracellular pH 6.0 ($n=4$). At pH 7.4, the small increase in inward current (measured at -100 mV) elicited by bath sulfate addition (from $-122 \pm 16 \text{ nA}$ to $-183 \pm 9 \text{ nA}$, $n=4$, $p < 0.05$) was unaccompanied by significant change in reversal potential. Taken together, the data suggest that Rv1739c function is expressed at very low level, if at all, in *Xenopus* oocytes.

3.4. Pharmacological properties of Rv1739c-associated sulfate uptake

As shown in Fig. 4A and B, sulfate uptake associated with induction of Rv1739c expression was abolished by the nonspecific inhibitors of the F1/F0 H^+ -ATPase, the sulfhydryl alkylator N-ethylmaleimide (NEM, 1 mM) and the oxidative phosphorylation uncoupler, carbonyl cyanide 3-chloro-phenyl-hydrazone (CCCP, 50 μM). Rv1739c-associated sulfate uptake was also abolished or nearly so by 5 mM of the sulfate analogs sulfite (Fig. 4C), selenate (Fig. 4D), and thiosulfate (Fig. 4E). But none of these inhibitors discriminated between native sulfate uptake and sulfate uptake attributable to Rv1739c expression. 5 mM of the sulfate analog molybdate, and the general anion transport inhibitors niflumate (100 μM , Fig. 4E) and di-isothiocyanato-4',4''-disulfono-stilbene disulfonate (DIDS, 1 mM, Fig. 4F) lowered induced sulfate uptake to uninduced levels. However, as 1 mM DIDS also nearly completely inhibited sulfate uptake by uninduced bacteria ($n=14$, not shown), it provided at this concentration little discrimination between native and Rv1739c-associated sulfate

uptake. The anion transport inhibitor NS3623 (100 μM) did not inhibit uninduced or induced sulfate uptake ($n=2$, not shown).

3.5. The transmembrane domain of Rv1739c is required for the increased sulfate uptake associated with induced overexpression

Rv1739c consists of two principal structural domains, the N-terminal transmembrane domain with a short predicted juxtamembrane cytoplasmic stretch, encompassed by aa 1–436, and the C-terminal cytoplasmic domain (aa 437–560) encompassing the “sulfate transporter and anti-sigma factor antagonist” (STAS) domain. Overexpression of 6His-tagged aa 1–436 transmembrane domain was associated with increased sulfate transport (Fig. 5A), whereas overexpression of the C-terminal aa 437–560 cytoplasmic STAS domain did not enhance sulfate uptake. This result is consistent with the membrane domain either mediating or being necessary for transport mediated by an associated native polypeptide. However, expression of the just the STAS domain itself, presented as aa 437–560, was not sufficient to support increased sulfate uptake by native transport proteins.

3.6. Rv1739c cysteine residues are not required for Rv1739c-associated increased sulfate uptake

Complete inhibition of sulfate uptake by NEM suggested that one or more of the three cysteine residues of Rv1739c might be essential for Rv1739c-associated sulfate uptake. Creation of Cys-less Rv1739c would provide not only a test of this hypothesis, but would also represent a valuable substrate




Fig. 5. A. ^{35}S -sulfate uptake is increased by induction of expression of the Rv1739c transmembrane domain (aa 1–436). B. ^{35}S -sulfate uptake is not increased by induction of expression of the C-terminal cytoplasmic domain of Rv1739c (aa 437–560) including its STAS domain. Number of measurements is within parentheses. *, $p < 0.05$.




Fig. 6. A. Schematic of the Cys-less Rv1739c polypeptide with its three cysteine-to-serine substitutions indicated. B. Induction of ^{35}S -sulfate uptake in cells expressing Cys-less Rv1739c. C. Inhibition by NEM (1 mM) and by CCCP (50 μM) of ^{35}S -sulfate uptake into cells induced to express Cys-less Rv1739c. Number of measurements is within parentheses. *, $p < 0.05$.

for scanning cysteine accessibility mutagenesis. Therefore, the two cysteine residues of the transmembrane domain and the single cysteine residue of the STAS domain were mutagenized to serine (Fig. 6A). However mutagenesis of all three cysteine residues neither reduced the IPTG-induced increase in sulfate uptake (Fig. 6B) nor altered the complete inhibition of either uninduced or induced levels of sulfate uptake by NEM (Fig. 6C). Moreover, the pH-dependence of Cys-less Rv1739c was unchanged (pH 6.0 > pH 7.0 > pH 8.0 > pH 5.0, $n=2$, not shown).

The Cys-less Rv1739c provided an opportunity to reintroduce single cysteine residues into the polypeptide (Fig. 7A) for subsequent tests of targeted sulfhydryl modification with methanethiosulfonate (MTS) reagents. Similar cysteine scanning mutagenesis has defined residues important for the function or folding of the phylogenetically related sulfate transporters Sultr1;2 of *A. thaliana* (Shibagaki and Grossman, 2004, 2006) and SHST1 of the tropical legume *Stylosanthes hamata* (Howitt, 2005). The IPTG-induced increment in Rv1739c-associated sulfate uptake was preserved in the protein products of all mutations introduced into the Cys-less polypeptide (Fig. 7B). These mutants could therefore be used, on the hypothesis that Rv1739c was mediating sulfate influx, to test the topographical disposition and the requirement for sulfate uptake of individual residues. Fig. 7C shows that sulfate uptake associated with expression

of Cys-less Rv1739c mutant P54C, but not by the Cys-less parent protein, was modestly inhibited by 1 mM MTSES. 1 mM concentrations of MTSET inhibited neither Cys-less Rv1739c nor its P54C mutant. However, 10 mM MTSET and MTSES each inhibited Cys-less Rv1739c-associated sulfate uptake by 60% ($n=6$, not shown). Cys-less Rv1739c was also inhibited 50% by 100 μM MTSEA, and completely by 1 mM, and the P54C mutant showed roughly similar sensitivity ($n=4$, not shown). Thus, the effects of MTS reagents on sulfate uptake were not mediated directly through Rv1739c.

3.7. Constitutive expression fails to rescue sulfate auxotrophy in the BCG *cysA* auxotroph strain EW44

The *M. bovis cysA* gene product has been previously defined by a global genetic complementation screen in rich medium as the unique sulfate uptake pathway of *M. bovis* (Wooff et al., 2002). We tested the hypothesis that constitutive overexpression of candidate sulfate transporter Rv1739c might complement the sulfate auxotrophy of the EWP44 strain of *M. bovis* lacking functional *cysA*. However, heterologous expression of Rv1739c-6His did not rescue EWP44 sulfate auxotrophy, whereas complementation with the *cysA* gene did so (not shown). Moreover, constitutive Rv1739c expression inhibited growth of EWP44 even in the presence of methionine. Thus, expression of Rv1739c in this setting appeared toxic.

3.8. Density-dependent induction of native sulfate uptake and Rv1739c-associated increased sulfate uptake are both abolished in the absence of the CysA component of the ABC sulfate uptake pathway

We tested the hypothesis that expression of CysA in *E. coli* might be required for enhancement of sulfate uptake by overexpression of Rv1739c. This experiment was of additional interest in testing the role of CysA in the density-dependent endogenous sulfate uptake system of *E. coli*. The *E. coli* strain BW25113 (the parent strain for the global library of insertional single gene knockouts (Baba et al., 2006)) displays growth




Fig. 7. A. Schematic of single cysteine substitutions introduced into the Cys-less Rv1739c polypeptide. B. Increased ^{35}S -sulfate uptake into cells induced to express Cys-less Rv1739c polypeptide containing the indicated single reintroduced cysteine residues. C. Effects of MTSET and MTSES on sulfate uptake induced in Cys-less Rv1739 and its P54C variant. Number of measurements is indicated within parentheses.




Fig. 8. A. Cell density-dependent induction of ^{35}S -sulfate transport in *E. coli* strain BW25113, but not in the *cysA* deletion strain JD21556. B. Arabinose treatment does not increase ^{35}S -sulfate uptake into JD21556 bacteria expressing Rv1739c in plasmid pBAD33. Number of measurements is indicated within parentheses. *, $p < 0.05$.

density-dependent increase in sulfate uptake (Fig. 8A) similar to that observed in Tumor cells. This density-stimulated sulfate uptake was completely abrogated in the JD21556 strain derived from BW25113 and lacking a functional *cysA* gene (Fig. 8A). This result suggested the possibility that not only the basal sulfate uptake but also the Rv1739c-stimulated sulfate uptake might be mediated by the CysA pathway.

Since the JD21556 strain has very low background sulfate flux, we tested whether improved signal-to-noise might benefit the stimulation of sulfate uptake by expression of Rv1739c. Rv1739c-6His was subcloned into pBAD33, transformed into JD21556, and incubated in the absence or presence of 0.002% L-arabinose. As shown in Fig. 8B, arabinose did not induce elevated expression of sulfate uptake in JD21556 cells harboring Rv1739c. Despite immunoblot evidence of Rv1739c induction (not shown), arabinose induction at 0.002% or (not shown) at concentrations from 100-fold lower to 100-fold higher) failed to increase Rv1739c-associated sulfate uptake in JD21556 cells. Thus, Rv1739c expression did not increase sulfate uptake in the absence of CysA expression.

4. Discussion

The large SLC26-related SulP family includes sulfate transporters in yeast and plants, but has been minimally characterized in bacteria. The only functional analysis of a bacterial SulP protein reported to date is that for the Na^+ -dependent HCO_3^-

transporter BicA of euryhaline cyanobacterial *Synechococcus* species (Price et al., 2004), a representative of the small class of SulP/CynT rosetta protein fusions of an SLC26-like anion transporter sequence with a type β -carbonic anhydrase sequence (Felce and Saier, 2004). The transport function of polypeptides of the more abundant SulP class characterized by presence of a C-terminal cytoplasmic STAS domain has not been reported. In addition, no bacterial or yeast SulP protein has been studied in isolated membranes or in reconstituted proteoliposomes.

Mycobacterial SulP transporters offer potential insight into the sulfur assimilation pathways leading to biosynthesis of sulfolipid pathogenicity determinants. The bacterial SulP polypeptides also provide attractive targets for structural analysis of protein folds homologous to those of the mammalian SLC26 anion transporter disease genes. Therefore we investigated the function of Rv1739c that, unique among the three examined recombinant SulP genes of the *M. tuberculosis* genome, accumulated to substantial levels after IPTG induction in Tuner DE3pLacI cells.

Although endogenous sulfate uptake in Tuner cells increases dramatically in late-log phase, Rv1739c overexpression in Tuner cells led to 2–5-fold increased sulfate uptake in early-to-mid log phase cells. This incremental sulfate uptake was maximal at pH 6 but abolished at pH 5, and was of high affinity with $K_{1/2}$ of 4 μ M. Rv1739c expression did not increase uptake of Cl^- , formate, or oxalate, and neither HCO_3^- nor nitrate inhibited sulfate uptake. Sulfate uptake was nearly abolished by the metabolic inhibitors NEM and CCCP, and the sulfate analogs sulfite, selenate, and thiosulfate. Sulfate uptake was inhibited to lesser extent by molybdate and the general anion transport inhibitors, niflumate and DIDS, but not by NS3623. Expression of the Rv1739c transmembrane domain was sufficient to increase sulfate uptake, but expression of the isolated C-terminal cytoplasmic domain did not affect sulfate uptake. Rv1739c-associated sulfate uptake was specific, insofar as no increase in sulfate uptake was associated with equivalent expression levels of *E. coli* SulP polypeptide ychM.

4.1. Role of the STAS domain in Rv1739c function

The function of the Rv1739c STAS domain remains unknown. A clue resides in the preliminary report that the *A. thaliana* Sultr1;2 STAS domain binds to the O-acetyl serine (thiol) lyase/cysteine synthase (OASTL) cytosolic isoform expressed in cells of the root cortex. This complex may constitute a sulfate assimilation metabolon (Shibagaki and Grossman, 2007), deletion or mutation of which increases resistance to selenate toxicity (El Kassis et al., 2007). Interestingly, *E. coli* CysA also can bind to OASTL/CysK (Arifuzzaman et al., 2006). STAS domains in plant SulP and mammalian SLC26 transporters are essential for plasma membrane targeting and for transport function. In SLC26A3, the STAS domain binds the R domain of the cystic fibrosis transmembrane regulator (Ko et al., 2004). Carbonic anhydrase II binding by SLC26A6 STAS domain is associated with increased SLC26A6-associated $\text{Cl}^-/\text{HCO}_3^-$ exchange activity (Alvarez et al., 2005).

The three cysteine residues of Rv1739c were non-essential for increased sulfate uptake. This finding resembles results reported for the SHST1 SulP sulfate transporter of a tropical legume (Howitt, 2005). However, unlike the STAS domain requirement for sulfate transport by *A. thaliana* Sultr1;2 (Shibagaki and Grossman, 2004), the STAS domain was dispensable for Rv1739c-associated enhancement of sulfate uptake. Moreover, modification of selected conserved residues homologous to those found important for sulfate transport by the plant SulP polypeptides SHST1 (Khurana et al., 2000; Shelden et al., 2003) and Sultr1;2 polypeptides (Shibagaki and Grossman, 2006) also appeared unimportant for Rv1739c-associated sulfate uptake. Expression of the Rv1739c STAS domain in the absence of its transmembrane domain did not enhance sulfate uptake.

4.2. Relation of the Rv1739c polypeptide to other SulP polypeptides

Rv1739c shares only 25% amino acid identity with the STAS domain-containing *M. tuberculosis* SulP polypeptide Rv1707, and only 23% identity with the cynT domain-containing *M. tuberculosis* SulP polypeptide Rv3273, a putative HCO_3^- transporter. Rv3273 polypeptide has been detected on Coomassie Blue G250-stained SDS-PAGE gels of alkali-extracted inner membrane fraction from *M. tuberculosis* (Mattow et al., 2007), and is present in *M. bovis* BCG membranes at 10-fold greater abundance (Schmidt et al., 2004). In contrast, the two STAS-domain-containing SulP proteins Rv1739c and Rv1707 have not been detected in proteomic screens of *M. tuberculosis*.

Rv1739c has a predicted pI 10.39, with basic residues distributed throughout both transmembrane and cytoplasmic domains. The distantly related SulP homolog Rv1707 also has a pI of 10.24. These alkaline pI values are rare among the *M. tuberculosis* proteome, in which the majority of polypeptides is highly acidic (Mattow et al., 2003). Rv1739c is marked by a 6 residue internal direct repeat, AVLAAT at aa 105–110 and again at 213–218, with AAVLA also at aa 90–94. The significance of these sequences beyond their hydrophobicity is unknown. Rv1739c from *M. tuberculosis* Strain H37Rv differs in the single residue R134 from the L134 present in strains CDC1551, F11, and c, as well as in the *M. bovis* ortholog. The smaller genome of *M. leprae* lacks all three SulP genes. The closest non-mycobacterial homolog of Rv1739c detected by BLAST is SulP polypeptide NP_600277 from *Corynebacterium glutamicum*, with 51% amino acid sequence identity.

The *P. aeruginosa* SulP polypeptide PA1647 shares 41% amino acid identity with *M. tuberculosis* Rv1739c. PA1647 mRNA abundance increased 3-fold upon sulfate deprivation, the same degree of increase as exhibited by the *P. aeruginosa* sulfate permease gene products CysT, CysW, and CysA. Moreover, PA1647 mRNA was increased 6-fold by growth of the cystic fibrosis *P. aeruginosa* isolate E601 on (sulfate-rich) mucin in sulfate-free conditions (Tralau et al., 2007). Sulfate starvation experiments revealed no upregulation of the ychM gene in *E. coli* (Gyaneshwar et al., 2005) or of any SulP gene

product in *B. subtilis* (Auger et al., 2002), but have not been published for *M. tuberculosis*. Nonetheless, the *P. aeruginosa* data provide evidence for a role of a Rv1739c-related polypeptide in the sulfur assimilation response, and suggest a direct or supporting role in sulfate transport function.

M. tuberculosis Rv1739c shares 23% amino acid sequence identity and 44% homology (NCBI BLAST) with trout *Onchorhynchus mykiss* Slc26a1 polypeptide (Katoh et al., 2006) across their aligned putative transmembrane domains and juxtamembrane regions, and 24% identity (45% homology) with eel *Anguilla japonica* kidney Slc26a1 (Nakada et al., 2005).

4.3. Regulation of Rv1739c gene expression

Several reports have demonstrated control of *M. tuberculosis* Rv1739c expression by environmental stressors believed to mimic the stresses of the host environment (Murphy and Brown, 2007). Thus, 24 h after *M. tuberculosis* infection of interferon- γ -activated macrophages, Rv1739c and Rv1707 mRNAs increased 2.5 and 4-fold, respectively. Whereas the increased expression of Rv1739c was independent of macrophage inducible nitric oxide synthase (iNOS) activity, induction of Rv1707 required iNOS activity (Schnappinger et al., 2003). An *in vitro* study noted 4-fold upregulation of Rv1739c mRNA in response to imposition of hypoxia in early log-phase liquid culture (Sherman et al., 2001). If in hypoxic conditions the ATP required for the ABC sulfate permease activity is limiting, an ATP-independent sulfate transporter might confer advantage. Although exposure in liquid culture to nitric oxide donors at high concentration led to 20-fold upregulation of Rv1739c mRNA levels (Ohno et al., 2003), lower concentrations believed better to model a "dormancy response" did not increase Rv1739c mRNA (Voskuil et al., 2003). *E. coli* SulP ychM mRNA levels have been shown to be upregulated 2–3 fold in glass wool biofilms (Ren et al., 2004).

4.4. Is the *M. tuberculosis* SulP polypeptide Rv1739c a sulfate transporter?

Both Tuner cells and the *E. coli* strain BW25113 exhibited strong induction of sulfate uptake in late-log and saturation phase of growth in liquid culture (Figs. 2 and 8). In the JD21556 derivative of BW25113 lacking CysA, both basal and density-induced sulfate uptake was almost completely abrogated (Fig. 8). Thus, CysA-mediated sulfate uptake in *E. coli* is strongly regulated by cell density in rich medium. This density-dependence may in part reflect sulfate depletion from the medium, since sulfate starvation of *E. coli* altered expression of the genes encoding the ABC sulfate permease subunits (Gyaneshwar et al., 2005).

Our experiments in rich medium showed that Rv1739c does not complement sulfate auxotrophy in the EWP44 BCG mutant deficient in CysA. However, interpretation of these experiments is confounded by the toxicity of constitutive Rv1739c expression in BCG strain EWP44. Similar toxicity was also reported for expression of the GFP-fusion protein of *E. coli* SulP polypeptide

ychM (<http://ecoli.aist.nara.ac.jp/GB6/info.jsp?id=JW5189>). The presence of CysA also proved necessary for Rv1739c-associated increased sulfate uptake in *E. coli*. Initial attempts to demonstrate intrinsic sulfate transport function of Rv1739c through sulfate flux studies in rightside-out vesicles prepared from IPTG-induced Tuner cells were unsuccessful. Thus, we conclude from our data that overexpression of Rv1739c in *E. coli* increases cellular sulfate uptake in a manner dependent upon expression (and presumed function) of CysA, the cytoplasmic ATP-binding subunit of the ABC sulfate permease. Our results suggest that Rv1739c may be a sulfate transporter that acts in conjunction with the CysTWA ABC sulfate permease.

If Rv1739c is indeed a sulfate transporter, its activity optimum at pH 6.0 is potentially consistent with the H⁺/sulfate cotransport mechanism proposed for plant (Buchner et al., 2004) and yeast SulP transporters, but the complete inhibition of Rv1739c-associated sulfate uptake at pH 5 remains unexplained. The lack of inhibition by extracellular HCO₃⁻ of Rv1739c overexpression-associated sulfate uptake into *E. coli* does not support a sulfate/bicarbonate exchange mechanism such as proposed for vertebrate Slc26a1 (Xie et al., 2002; Markovich and Aronson, 2007) and consistent with carbonic anhydrase-dependence of sulfate excretion by teleost kidney (Renfro et al., 1999; Nakada et al., 2005; Pelis et al., 2005; Katoh et al., 2006). In addition, the lack of upregulated oxalate transport by Rv1739c overexpression and lack of inhibition by oxalate of associated sulfate uptake both suggest a substrate specificity distinct from that of oxalate-transporting mouse Slc26a1 (Xie et al., 2002).

Wooff et al. (2002) showed that the CysTWA sulfate permease system was the only essential sulfate transporter of BCG in rich medium and in amino acid-deprived medium. However, deletion of the *cysA* gene did not impair BCG survival and proliferation in normal mice. Thus, it remains possible that Rv1739c and/or other SulP proteins of mycobacteria play important roles mediating or regulating sulfate transport during host infection, a condition not optimally modeled by standard *in vitro* culture. Sulfate uptake mediated by cystTWA and Rv1739c polypeptides may direct sulfate into different metabolic pathways built around distinct enzymatic assemblies that have been described as "enzoskeletons" (Norris et al., 1996).

4.5. Conclusion

The SLC26-related SulP polypeptide Rv1739c of *M. tuberculosis* has been overexpressed in *E. coli* and shown to increase sulfate uptake into intact bacteria. The increased sulfate uptake occurs by a mechanism requiring the cytoplasmic CysA subunit of the ABC sulfate permease. The transmembrane domain of Rv1739c suffices for this activity. Further experiments will be needed to test intrinsic sulfate transport activity of isolated Rv1739c polypeptide.

Acknowledgements

This work was supported by NIH grants R01 DK43495 (SLA), DK34854 (Harvard Digestive Diseases Center funds to SLA) and T32 DK07199 (Beth Israel Deaconess Renal Training Grant to JSC). We thank Drs. E. Wooff and P. Wheeler

(Veterinary Laboratories Agency, Surrey, UK) for the kind gifts of *M. bovis* strain EWP44 and *M. bovis* expression plasmid pSM-101EW.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.cbpa.2007.12.005](https://doi.org/10.1016/j.cbpa.2007.12.005).

References

- Alper, S.L., 2006. Molecular physiology of SLC4 anion exchangers. *Exp. Physiol.* 91, 153–161.
- Alvarez, B.V., Vilas, G.L., Casey, J.R., 2005. Metabolism disruption: a mechanism that regulates bicarbonate transport. *Embo J.* 24, 2499–2511.
- Arifuzzaman, M., Maeda, M., Itoh, A., Nishikata, K., Takita, C., Saito, R., Ara, T., Nakahigashi, K., Huang, H.C., Hirai, A., Tsuzuki, K., Nakamura, S., Altaf-Ul-Amin, M., Oshima, T., Baba, T., Yamamoto, N., Kawamura, T., Ioka-Nakamichi, T., Kitagawa, M., Tomita, M., Kanaya, S., Wada, C., Mori, H., 2006. Large-scale identification of protein-protein interaction of *Escherichia coli* K-12. *Genome Res.* 16, 686–691.
- Auger, S., Danchin, A., Martin-Verstraete, I., 2002. Global expression profile of *Bacillus subtilis* grown in the presence of sulfate or methionine. *J. Bacteriol.* 184, 5179–5186.
- Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M., Wanner, B.L., Mori, H., 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* 2, 2006 0008.
- Buchner, P., Takahashi, H., Hawkesford, M.J., 2004. Plant sulphate transporters: co-ordination of uptake, intracellular and long-distance transport. *J. Exp. Bot.* 55, 1765–1773.
- Casula, S., Shmukler, B.E., Wilhelm, S., Stuart-Tilley, A.K., Su, W., Chernova, M.N., Brugnara, C., Alper, S.L., 2001. A dominant negative mutant of the KCC1 K-Cl cotransporter: both N- and C-terminal cytoplasmic domains are required for K-Cl cotransport activity. *J. Biol. Chem.* 276, 41870–41878.
- Chernova, M.N., Jiang, L., Friedman, D.J., Darman, R.B., Lohi, H., Kere, J., Vanderpe, D.H., Alper, S.L., 2005. Functional comparison of mouse *slc26a6* anion exchanger with human SLC26A6 polypeptide variants: differences in anion selectivity, regulation, and electrogenicity. *J. Biol. Chem.* 280, 8564–8580.
- Clark, J.S., Vanderpe, D.H., Chernova, M.N., Heneghan, J.F., Stewart, A.K., Alper, S.L., in press. Species differences in Cl⁻ affinity and in electrogenicity of SLC26A6-mediated oxalate/Cl⁻ exchange correlate with the distinct human and mouse susceptibilities to nephrolithiasis. *J. Physiol.*
- El Kassis, E., Cathala, N., Rouached, H., Fourcroy, P., Berthomieu, P., Terry, N., Davidian, J.C., 2007. Characterization of a selenate-resistant *Arabidopsis* mutant. Root growth as a potential target for selenate toxicity. *Plant Physiol.* 143, 1231–1241.
- Fan, F., Luxenburger, A., Painter, G.F., Blanchard, J.S., 2007. Steady-state and pre-steady-state kinetic analysis of *Mycobacterium smegmatis* cysteine ligase (MshC). *Biochemistry* 46, 11421–11429.
- Felce, J., Saier Jr., M.H., 2004. Carbonic anhydrases fused to anion transporters of the SulP family: evidence for a novel type of bicarbonate transporter. *J. Mol. Microbiol. Biotechnol.* 8, 169–176.
- Guzman, L.M., Belin, D., Carson, M.J., Beckwith, J., 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J. Bacteriol.* 177, 4121–4130.
- Gyaneshwar, P., Paliy, O., McAuliffe, J., Popham, D.L., Jordan, M.I., Kustu, S., 2005. Sulfur and nitrogen limitation in *Escherichia coli* K-12: specific homeostatic responses. *J. Bacteriol.* 187, 1074–1090.
- Howitt, S.M., 2005. The role of cysteine residues in the sulphate transporter, SHST1: construction of a functional cysteine-less transporter. *Biochim. Biophys. Acta* 1669, 95–100.
- Katoh, F., Tresguerres, M., Lee, K.M., Kaneko, T., Aida, K., Goss, G.G., 2006. Cloning of rainbow trout SLC26A1: involvement in renal sulfate secretion. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 290, R1468–R1478.
- Kere, J., 2006. Overview of the SLC26 family and associated diseases. *Novartis Found. Symp.* 273, 2–11 discussion 11–18, 261–264.
- Khurana, O.K., Coupland, L.A., Shelden, M.C., Howitt, S.M., 2000. Homologous mutations in two diverse sulphate transporters have similar effects. *FEBS Lett.* 477, 118–122.
- Ko, S.B., Zeng, W., Dorwart, M.R., Luo, X., Kim, K.H., Millen, L., Goto, H., Naruse, S., Soyombo, A., Thomas, P.J., Muallem, S., 2004. Gating of CFTR by the STAS domain of SLC26 transporters. *Nat. Cell Biol.* 6, 343–350.
- Kurschat, C.E., Shmukler, B.E., Jiang, L., Wilhelm, S., Kim, E.H., Chernova, M.N., Kinne, R.K., Stewart, A.K., Alper, S.L., 2006. Alkaline-shifted pHo sensitivity of AE2c1-mediated anion exchange reveals novel regulatory determinants in the AE2 N-terminal cytoplasmic domain. *J. Biol. Chem.* 281, 1885–1896.
- Lecanda, J., Urtasun, R., Medina, J.F., 2000. Molecular cloning and genomic organization of the mouse AE2 anion exchanger gene. *Biochem. Biophys. Res. Commun.* 276, 117–124.
- Lindblow-Kull, C., Kull, F.J., Shrift, A., 1985. Single transporter for sulfate, selenate, and selenite in *Escherichia coli* K-12. *J. Bacteriol.* 163, 1267–1269.
- Markovich, D., Aronson, P.S., 2007. Specificity and regulation of renal sulfate transporters. *Annu. Rev. Physiol.* 69, 361–375.
- Mattow, J., Schaible, U.E., Schmidt, F., Hagens, K., Siejak, F., Brestrich, G., Haeselbarth, G., Muller, E.C., Jungblut, P.R., Kaufmann, S.H., 2003. Comparative proteome analysis of culture supernatant proteins from virulent *Mycobacterium tuberculosis* H37Rv and attenuated *M. bovis* BCG Copenhagen. *Electrophoresis* 24, 3405–3420.
- Mattow, J., Siejak, F., Hagens, K., Schmidt, F., Koehler, C., Treumann, A., Schaible, U.E., Kaufmann, S.H., 2007. An improved strategy for selective and efficient enrichment of integral plasma membrane proteins of mycobacteria. *Proteomics* 7, 1687–1701.
- Mount, D.B., Romero, M.F., 2004. The SLC26 gene family of multifunctional anion exchangers. *Pflugers Arch.* 447, 710–721.
- Murphy, D.J., Brown, J.R., 2007. Identification of gene targets against dormant phase *Mycobacterium tuberculosis* infections. *BMC Infect. Dis.* 7, 84.
- Nakada, T., Zandi-Nejad, K., Kurita, Y., Kudo, H., Broumand, V., Kwon, C.Y., Mercado, A., Mount, D.B., Hirose, S., 2005. Roles of Slc13a1 and Slc26a1 sulfate transporters of eel kidney in sulfate homeostasis and osmoregulation in freshwater. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 289, R575–R585.
- Norris, V., Turnock, G., Sigee, D., 1996. The *Escherichia coli* zenoskeleton. *Mol. Microbiol.* 19, 197–204.
- Ohno, H., Zhu, G., Mohan, V.P., Chu, D., Kohno, S., Jacobs Jr., W.R., Chan, J., 2003. The effects of reactive nitrogen intermediates on gene expression in *Mycobacterium tuberculosis*. *Cell. Microbiol.* 5, 637–648.
- Pelis, R.M., Edwards, S.L., Kunigelis, S.C., Claiborne, J.B., Renfro, J.L., 2005. Stimulation of renal sulfate secretion by metabolic acidosis requires Na⁺/H⁺ exchange induction and carbonic anhydrase. *Am. J. Physiol. Renal Physiol.* 289, F208–F216.
- Price, G.D., Woodger, F.J., Badger, M.R., Howitt, S.M., Tucker, L., 2004. Identification of a SulP-type bicarbonate transporter in marine cyanobacteria. *Proc. Natl. Acad. Sci. U.S.A.* 101, 18228–18233.
- Quick, M., Wright, E.M., 2002. Employing *Escherichia coli* to functionally express, purify, and characterize a human transporter. *Proc. Natl. Acad. Sci. U.S.A.* 99, 8597–8601.
- Ren, D., Bedzyk, L.A., Thomas, S.M., Ye, R.W., Wood, T.K., 2004. Gene expression in *Escherichia coli* biofilms. *Appl. Microbiol. Biotechnol.* 64, 515–524.
- Renfro, J.L., Maren, T.H., Zeien, C., Swenson, E.R., 1999. Renal sulfate secretion is carbonic anhydrase dependent in a marine teleost, *Pleuronectes americanus*. *Am. J. Physiol.* 276, F288–F294.
- Romero, M.F., Fulton, C.M., Boron, W.F., 2004. The SLC4 family of HCO₃⁻ transporters. *Pflugers Arch.* 447, 495–509.
- Rosenberg, N.K., Lee, R.W., Yancey, P.H., 2006. High contents of hypotaurine and thiotaurine in hydrothermal-vent gastropods without thiotrophic endosymbionts. *J. Exp. Zool. A Comp. Exp. Biol.* 305, 655–662.
- Schelle, M.W., Bertozzi, C.R., 2006. Sulfate metabolism in mycobacteria. *ChemBiochem* 7, 1516–1524.
- Schmidt, F., Donahoe, S., Hagens, K., Mattow, J., Schaible, U.E., Kaufmann, S.H., Aebbersold, R., Jungblut, P.R., 2004. Complementary analysis of the *Mycobacterium tuberculosis* proteome by two-dimensional electrophoresis and isotope-coded affinity tag technology. *Mol. Cell. Proteomics* 3, 24–42.
- Schnappinger, D., Ehrt, S., Voskuil, M.I., Liu, Y., Mangan, J.A., Monahan, I.M., Dolganov, G., Efron, B., Butcher, P.D., Nathan, C., Schoolnik, G.K., 2003.

- Transcriptional adaptation of *Mycobacterium tuberculosis* within macrophages: insights into the phagosomal environment. *J. Exp. Med.* 198, 693–704.
- Shelden, M.C., Loughlin, P., Tierney, M.L., Howitt, S.M., 2003. Interactions between charged amino acid residues within transmembrane helices in the sulfate transporter SHST1. *Biochemistry* 42, 12941–12949.
- Sherman, D.R., Voskuil, M., Schnappinger, D., Liao, R., Harrell, M.I., Schoolnik, G.K., 2001. Regulation of the *Mycobacterium tuberculosis* hypoxic response gene encoding alpha-crystallin. *Proc. Natl. Acad. Sci. U.S.A.* 98, 7534–7539.
- Shibagaki, N., Grossman, A.R., 2004. Probing the function of STAS domains of the Arabidopsis sulfate transporters. *J. Biol. Chem.* 279, 30791–30799.
- Shibagaki, N., Grossman, A.R., 2006. The role of the STAS domain in the function and biogenesis of a sulfate transporter as probed by random mutagenesis. *J. Biol. Chem.* 281, 22964–22973.
- Shibagaki, N., Grossman, A.R., 2007. Functions of the STAS domain in the Arabidopsis sulfate transporter, Sultr1;2. *Botany and Plant Biology 2007 Joint Congress*. p. Abst. 1462 poster P11018.
- Tralau, T., Vuilleumier, S., Thibault, C., Campbell, B.J., Hart, C.A., Kertesz, M.A., 2007. Transcriptomic analysis of the sulfate starvation response of *Pseudomonas aeruginosa*. *J. Bacteriol.* 189, 6743–6750.
- Voskuil, M.I., Schnappinger, D., Visconti, K.C., Harrell, M.I., Dolganov, G.M., Sherman, D.R., Schoolnik, G.K., 2003. Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *J. Exp. Med.* 198, 705–713.
- Wooff, E., Michell, S.L., Gordon, S.V., Chambers, M.A., Bardarov, S., Jacobs Jr., W.R., Hewinson, R.G., Wheeler, P.R., 2002. Functional genomics reveals the sole sulphate transporter of the *Mycobacterium tuberculosis* complex and its relevance to the acquisition of sulphur in vivo. *Mol. Microbiol.* 43, 653–663.
- Xie, Q., Welch, R., Mercado, A., Romero, M.F., Mount, D.B., 2002. Molecular characterization of the murine Slc26a6 anion exchanger: functional comparison with Slc26a1. *Am. J. Physiol. Renal. Physiol.* 283, F826–F838.
- Yoshimoto, N., Inoue, E., Watanabe-Takahashi, A., Saito, K., Takahashi, H., 2007. Posttranscriptional regulation of high-affinity sulfate transporters in Arabidopsis by sulfur nutrition. *Plant Physiol.* 145, 378–388.