



Available online at www.sciencedirect.com



Fisheries Research 75 (2005) 204–205

**FISHERIES
RESEARCH**

www.elsevier.com/locate/fishres

Erratum

Erratum to “Using a modified Nordmøre grid for by-catch reduction in the Portuguese crustacean trawl fishery” [Fish. Res. 71 (2005) 223–239]

Paulo Fonseca ^{a,*}, Aida Campos ^a, Roger B. Larsen ^b, Teresa C. Borges ^c, Karim Erzini ^c

^a INIAP/IPIMAR—Portuguese Institute for Agriculture and Fisheries Research, Avenida de Brasília, 1449-006 Lisboa, Portugal

^b The Norwegian College of Fishery Science, University of Tromsø, N-9037 Tromsø, Norway

^c Centre of Marine Sciences (CCMAR), University of the Algarve, Campus de Gambelas, 8000-117 Faro, Portugal

The Publisher regrets that an error occurred in Fig. 6 of the above article, which is corrected as follows. The Publisher apologises for any inconvenience caused.

DOI of original article:10.1016/j.fishres.2004.08.018.

* Corresponding author. Tel.: +351 213 027 163; fax: +351 213 015 948.

E-mail addresses: pfonseca@ipimar.pt (P. Fonseca), acampos@ipimar.pt (A. Campos), rogerl@nfh.uit.no (R.B. Larsen), tborges@ualg.pt (T.C. Borges), kerzini@ualg.pt (K. Erzini).




Fig. 6. Length frequency distributions for the target species and blue whiting in 2001 (a–d) and 2002 (e–h). Thin line: total catch; thick line: inner codend; dashed line: codend; dashed line with circles: cover.