

available at www.sciencedirect.comwww.elsevier.com/locate/brainres**BRAIN
RESEARCH**

Research Report

Modulation of spontaneous hippocampal synaptic events with 5-hydroxyindole, 4OH-GTS-21, and rAAV-mediated $\alpha 7$ nicotinic receptor gene transfer

Jeffrey S. Thinschmidt, Gretchen Y. López-Hernández, Ke Ren, Michael A. King, Edwin M. Meyer, Roger L. Papke*

Department of Pharmacology and Therapeutics, University of Florida, Gainesville, Florida 32610, USA

ARTICLE INFO

Article history:

Accepted 1 February 2008

Available online 14 February 2008

Keywords:

 $\alpha 7$

Gene transfer

Nicotinic

rAAV

Hippocampus

5-hydroxyindole (5-HI)

GTS-21

DMXB

ABSTRACT

One approach to treatment of negative cognitive effects associated with Alzheimer's disease and schizophrenia may involve activation of neuronal $\alpha 7$ nicotinic acetylcholine receptors (nAChRs). We used the $\alpha 7$ -selective partial agonist 3-(4-hydroxy, 2-methoxybenzylidene)anabaseine (4OH-GTS-21), the $\alpha 7$ modulator 5-hydroxyindole (5-HI), and recombinant adeno-associated virus (rAAV)-mediated $\alpha 7$ gene transfer in order to test the hypothesis whether combining these strategies would significantly increase indirect measures of $\alpha 7$ nAChR function, including measures of spontaneous synaptic events in CA1 pyramidal cells. 5-HI (1 mM), and 5-HI (1 mM)+4OH-GTS-21 (5 μ M) increased the frequency of APV- and NBQX-sensitive currents, while 5-HI+4OH-GTS-21 increased the frequency and amplitude of bicuculline-sensitive currents. Effects on EPSCs were blocked with tetrodotoxin (TTX) (1 μ M), but not by methyllycaconitine (MLA) (50 nM). Neither TTX nor MLA reduced the potentiation of IPSC frequencies. However, TTX blocked, and in some cases MLA reduced, the potentiation of IPSC amplitudes. These data suggest that effects of 5-HI+4OH-GTS-21 on EPSC frequency were associated with action potential-dependent transmitter release produced by 5HI, and that potentiation of IPSC amplitudes resulted at least in part, from activation of $\alpha 7$ nAChRs. Finally, rAAV-mediated $\alpha 7$ gene transfer did not alter the magnitude of effects produced by 5-HI or 5-HI+4OH-GTS-21. Thus, although we previously showed that direct measures of $\alpha 7$ nAChR function were enhanced by $\alpha 7$ gene transfer, indirect measures of $\alpha 7$ nAChRs function were not significantly enhanced by combining $\alpha 7$ gene transfer with either agonist activation or positive allosteric modulation of $\alpha 7$ nAChRs.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

Nicotinic acetylcholine receptors (nAChRs) containing the $\alpha 7$ subunit are expressed at high levels in the hippocampus.

Unlike the high affinity nicotine-binding receptors of the brain, they can be activated by both acetylcholine and its precursor choline, but they display rapid desensitization to high concentrations of agonist, and are selectively blocked by methyl-

* Corresponding author. Department of Pharmacology and Therapeutics, 100267 JHMHC, 1600 SW Archer Road, University of Florida, College of Medicine, Gainesville, FL 32610, USA. Fax: +1 352 392 9696.

E-mail address: rlpapke@ufl.edu (R.L. Papke).

lycaconitine (MLA) and α -bungarotoxin. Responses mediated by $\alpha 7$ receptors can be readily detected by agonist applications to the cell bodies of hippocampal interneurons, and indirect measurements have suggested that they also function in the dendrites of hippocampal pyramidal cells (Ji and Dani, 2000). Disruption of $\alpha 7$ nAChR function has been implicated in Alzheimer's disease (AD) and schizophrenia (Court et al., 1999; Freedman et al., 1997; Guan et al., 2000), leading to their recognition as potentially important therapeutic targets for the treatment of these conditions. Because of their high permeability to calcium (Seguela et al., 1993), it is likely that $\alpha 7$ nAChRs modulate the release of many neurotransmitters at a variety of synapses. Activation of $\alpha 7$ nAChRs can modulate the release of GABA (Alkondon et al., 1997; Wanaverbecq et al., 2007), glutamate (Gray et al., 1996; McGehee et al., 1995), dopamine (Schilstrom et al., 1998), and noradrenaline (Li et al., 1998). Also, it has been shown that $\alpha 7$ nAChRs regulate the

excitability of CA1 pyramidal cells (Dani, 2000) and dentate granule cells (Frazier et al., 2003).

Several strategies are available for increasing $\alpha 7$ nAChR function including agonist activation, allosteric modulation, and $\alpha 7$ nAChR gene delivery. Agonist activation of $\alpha 7$ nAChRs has been demonstrated with the compound 3-(2,4 dimethoxybenzylidene)anabaseine (GTS-21 or DMXB), a selective partial agonist for rat $\alpha 7$ nAChRs. GTS-21 improves memory-related behaviors in aged rats (Arendash et al., 1995), aged rabbits (Woodruff-Pak et al., 1994), nonhuman primates (Briggs et al., 1997), and nucleus-basalis lesioned rats (Meyer et al., 1998a). Although GTS-21 has a lower efficacy for human $\alpha 7$ nAChRs than for rat $\alpha 7$ nAChR, its primary human metabolite, 3-(4-hydroxy, 2-methoxy-benzylidene)anabaseine (4OH-GTS-21) exhibits greater efficacy than GTS-21 for both rat and human $\alpha 7$ nAChRs, and compared to GTS-21, produces a better equilibrium between activation and agonist-dependent




Fig. 1 – Excitatory spontaneous synaptic currents in CA1 pyramidal cells at -60 mV and inhibitory synaptic currents at 0 mV are blocked by NBQX+APV and bicuculline respectively. The scatter plots in A and B were derived from averaging the number of events in 30 s intervals (\pm SEM). Baseline rates were recorded for 2 min (120 s) before bath application of the specified antagonist (thick black bars). **A:** Scatter plot (left) showing group means \pm SEM ($n=5$) and recordings from individual cells before (1) and following bath application of 10 mM NBQX and 40 mM APV (2) (note the absence of events shown in 2). **B:** Scatter plot (left) showing group means \pm SEM ($n=5$) and recordings from individual cells before (3) and following bath application of 30 mM bicuculline (4) (note the absence of events shown in 4). Scale bars: (1 and 2: 10 pA and 2.5 s) and (3 and 4: 20 pA and 2.5 s.). Any apparent differences in time course of the bath applied drugs are likely to reflect the relative potencies of the specific agents and their ability to penetrate the tissue.

inhibition and/or desensitization (Meyer et al., 1998b). Consequently, 4OH-GTS-21 may cause more receptor activation over extended intervals than other agonists which may induce greater inhibition. Also, like GTS-21, 4OH-GTS-21 produces positive cognitive and cytoprotective effects in rats (Ren et al., 2007a) and 4OH-GTS-21 has been shown to be able to increase spontaneous firing rates in $\alpha 7$ -expressing neurons (Uteshev et al., 2003).

Agonist activation of $\alpha 7$ receptors may be complicated in the treatment of AD by several factors. For example, there are reductions in hippocampal cholinergic innervation (Mesulam, 2004; Mufson et al., 2003), and $\alpha 7$ receptor function is reduced in animal models of cholinergic hypofunction (Thinschmidt et al., 2005). Also, there is an increase in astrocytic neurotransmitter receptor expression in AD (Teaktong et al., 2003). Thus, it may be practical to pursue alternative treatments such as gene therapy for AD. Accordingly, we recently reported (Ren et al., 2007b) a technique to elevate $\alpha 7$ nAChRs in a neuron-selective manner using recombinant adeno-associated virus (rAAV)-mediated delivery of $\alpha 7$ nAChR transgene directly into the hippocampus. We found that this resulted in an increase in [3 H]MLA binding in wild-type and $\alpha 7$ -receptor knockout (KO) mice, functional $\alpha 7$ receptors in $\alpha 7$ KO mice, substantial increases in the magnitude of ACh-evoked currents in stratum radiatum interneurons, and improved acquisition performance in the Morris water task (Ren et al., 2007b).

Known positive modulators of $\alpha 7$ nAChR include ivermectin (Krause et al., 1998), galantamine (Santos et al., 2002), PNU-120596 (Hurst et al., 2005), TQS (4,5,9b-tetrahydro-3-H-cyclopenta [c] quinoline-8-sulfonic acid amide) (Gronlien et al., 2007), and 5-hydroxyindole (5-HI) (Gurley et al., 2000; Zwart et al., 2002). However, 5-HI also produces a number of other physiological effects. For example, in the hippocampus 5-HI increases glutamate release, the amplitude of population spikes, and the amplitude of evoked excitatory and inhibitory postsynaptic potentials (Mannaioni et al., 2003). Also, 5-HI increases the frequency and amplitude of spontaneous GABAergic inhibitory postsynaptic currents (IPSCs) (Mannaioni et al., 2003; Mok and Kew, 2006). Thus, 5-HI is an effective positive allosteric modulator of $\alpha 7$ nAChRs that facilitates the output of both excitatory and inhibitory neurotransmission, although it is unclear from previous studies if all of the effects reported for 5-HI arise strictly from the positive modulation of $\alpha 7$ nAChRs.

In the present report we combined 4OH-GTS-21 treatment, expected to produce direct activation of $\alpha 7$ nAChRs, with allosteric modulation of $\alpha 7$ nAChRs using 5-HI, and rAAV-mediated $\alpha 7$ nAChR gene transfer in order to test the hypothesis that combining these strategies would produce increases in several potential sequella of increased $\alpha 7$ nAChR function, specifically the frequency and amplitude of spontaneous synaptic events in hippocampal CA1 pyramidal cells. We also tested the hypothesis that increasing the function and number of $\alpha 7$ nAChRs through $\alpha 7$ gene transfer would increase the magnitude of effects produced by bath application of 5-HI, 4OH-GTS-21, and 5-HI+4OH-GTS-21. Additionally, we compared the frequency and amplitude of spontaneous synaptic events in stratum radiatum interneurons from normal animals and animals that received $\alpha 7$ gene transfer.

2. Results

2.1. Pharmacological identification of spontaneous synaptic currents in CA1 pyramidal cells

In order to determine the pharmacological identity of spontaneous synaptic currents in CA1 pyramidal cells under our experimental conditions, we measured the effects of bath application of 10 μ M NBQX and 40 μ M 2-amino-5-phosphonopentanoic acid (APV) on inward currents recorded at -60 mV (putative excitatory postsynaptic currents (EPSCs)) using a K^+ based internal solution (Fig. 1A). The effects of 30 μ M bicuculline methiodide on outward currents were recorded at 0 mV (putative inhibitory postsynaptic currents (IPSCs)) using a Cs-based




Fig. 2 – Group data showing the effects of 4OH-GTS-21 (black bar), 5-HI (dark gray bar), 5-HI+4OH-GTS-21 (light gray bar), and 5-HI+4OH-GTS-21+TTX (hatched bar) on the frequency (A) and amplitude (B) of spontaneous EPSCs in hippocampal CA1 pyramidal cells from normal animals. * indicates statistically significant changes from baseline amplitude and/or frequency where $p < 0.05$.




Fig. 3 – Examples of the modulation of spontaneous synaptic events by 5-HI+4OH-GTS-21, which produced an increase in the frequency of spontaneous EPSCs in hippocampal pyramidal cells (A) and an increase the frequency and amplitude of spontaneous IPSCs in hippocampal pyramidal cells (B). Baseline synaptic events are shown on the left, and events in the presence of 5-HI+4OH-GTS-21 are shown on the right. In A the record was taken from a normal animal, and in B from an animal that received $\alpha 7$ gene transfer. Scale bars: A: (10 pA and 2.5 s) and B: (100 pA and 2.5 s).

internal solution (Fig. 1B). The spontaneous inward currents recorded under basal conditions were completely eliminated following bath application of NBQX and APV, confirming that these currents were EPSCs mediated by AMPA and/or NMDA receptors. Likewise, the putative spontaneous IPSCs recorded under basal conditions were completely eliminated following bath application of bicuculline methiodide, indicating that these currents were mediated by GABA-A receptors.

2.2. Drug effects on spontaneous EPSCs in CA1 pyramidal cells from normal animals

5-HI (1 mM) and 5-HI (1 mM)+4OH-GTS-21 (5 μ M) produced a significant increase in the frequency of EPSCs (Figs. 2A and 3A, Table 1). In contrast, when applied alone, 4OH-GTS-21 (5 μ M) had no significant effects on the frequency or amplitude of spontaneous EPSCs (Fig. 2A,B, Table 1). The increased frequency produced by 5-HI+4OH-GTS-21 was reduced by application of tetrodotoxin

(TTX) (1 μ M) (Fig. 2A). There was no effect of 5-HI+4OH-GTS-21 on the amplitude of EPSCs (Fig. 2B, Table 1), but EPSC amplitudes were significantly ($p < 0.01$) reduced when application of 5-HI+4OH-GTS-21 was followed by TTX (Fig. 2B, Table 1).

2.3. Drug effects on spontaneous EPSCs in CA1 pyramidal cells from animals that received $\alpha 7$ gene transfer

As seen in cells from normal animals, 5-HI (1 mM) and 5-HI (1 mM)+4OH-GTS-21 (5 μ M) both increased the frequency of EPSCs (Fig. 4A, Table 1) following $\alpha 7$ gene transfer. The effects produced by 5-HI were not reversed with bath application of methyllycaconitine (MLA) 50 nM, as shown in Fig. 4A and Table 1. When applied alone, 4OH-GTS-21 produced no significant effect on the frequency of EPSCs (Fig. 4A, Table 1), and there was no significant effect of 5-HI, 4OH-GTS-21, 5-HI+4OH-GTS-21, or 5-HI+MLA on the amplitudes of spontaneous EPSCs (Fig. 4B, Table 1).

2.4. Drug effects on spontaneous IPSCs in CA1 pyramidal cells from normal animals

When applied alone, 4OH-GTS-21 (5 μ M) produced no significant change in the frequency or amplitude of IPSCs (Fig. 5A,B, Table 2). In contrast, 5-HI (1 mM)+4OH-GTS-21 (5 μ M) increased the frequency and amplitude of IPSCs (Fig. 5A,B, Table 2). TTX (1 μ M) blocked and MLA reduced the increase in amplitude (Fig. 5B) but did not affect the increase in frequency (Fig. 5A). Although in the presence of 5-HI+4OH-GTS-21, the mean amplitude of IPSCs was 45.2 ± 7.9 pA, it should be noted that in some cases very large outward currents were recorded. These currents measured in excess of 1 nA but nevertheless were blocked with picrotoxin and bicuculline methiodide (data not shown). The increase in the frequency of IPSCs was not blocked by either TTX or MLA (Fig. 5A).

2.5. Drug effects on spontaneous IPSCs in CA1 pyramidal cells from animals that received $\alpha 7$ gene transfer

5-HI+4OH-GTS-21 produced a significant increase in the amplitude and frequency of IPSCs (Figs. 6A,B, and 3B, Table 2). These effects were not reversed with bath application of MLA (50 nM) (Fig. 6A,B). In contrast, when applied alone, 4OH-GTS-

Table 1 – EPSCs

Treatment	Baseline Hz	Treatment Hz	t-test	Baseline pA	Treatment pA	t-test	N
Normals							
4OH-GTS-21	0.21 \pm 0.04	0.27 \pm 0.06	–	15.2 \pm 0.7	14.2 \pm 1.0	–	9
5-HI	0.21 \pm 0.03	0.44 \pm 0.11	*	17.3 \pm 1.7	14.2 \pm 1.4	–	9
5-HI+4OH-GTS-21	0.21 \pm 0.04	0.62 \pm 0.17	*	15.2 \pm 0.7	15.5 \pm 1.8	–	9
5-HI+4OH-GTS-21+TTX	0.26 \pm 0.07	0.41 \pm 0.12	–	20.1 \pm 1.8	14.3 \pm 1.1	*	5
Gene transfer							
4OH-GTS-21	0.31 \pm 0.04	0.28 \pm 0.03	–	16.9 \pm 0.9	16.2 \pm 1.0	–	22
5-HI	0.48 \pm 0.02	1.0 \pm 0.22	*	17.4 \pm 1.8	18.6 \pm 1.6	–	8
5-HI+4OH-GTS-21	0.26 \pm 0.03	0.80 \pm 0.16	**	17.5 \pm 1.2	18.6 \pm 1.8	–	10
5-HI+MLA	0.39 \pm 0.44	0.84 \pm 0.28	–	16.6 \pm 1.8	15.8 \pm 1.3	–	4

Raw data showing the frequencies and amplitudes of EPSCs \pm SEM in CA1 pyramidal cells from normal animals and animals that received $\alpha 7$ gene transfer in baseline and drug treatment conditions. * $p < 0.05$, ** $p < 0.01$.




Fig. 4 – Group data showing the effects of 4OH-GTS-21 (black bar), 5-HI (gray bar), 5-HI + 4OH-GTS-21 (light gray bar), and 5-HI + MLA (white bar) on the frequency (A) and amplitude (B) of spontaneous EPSCs in hippocampal CA1 pyramidal cells from animals that received $\alpha 7$ gene transfer. * indicates statistically significant changes from baseline amplitude and/or frequency where $p < 0.05$ and ** where $p < 0.01$.

21 (5 μ M) produced no change in the frequency or amplitude of IPSCs (Fig. 6A,B, Table 2).

2.6. Comparison of drug effects on spontaneous synaptic events in CA1 pyramidal cells from normal animals and animals that received $\alpha 7$ gene transfer

The magnitude of increases in frequency and amplitude produced by bath application of 5-HI and 5-HI+4OH-GTS-21 reported above was not significantly different in cells from animals that received $\alpha 7$ gene transfer compared to cells from normal animals. Also, we found no significant difference in the baseline frequency or amplitude of EPSCs or IPSCs following $\alpha 7$ gene transfer, and no effects of MLA when applied alone (data not shown).

2.7. Comparison of the frequency and amplitude of spontaneous synaptic events in stratum radiatum interneurons from normal animals and animals that received $\alpha 7$ gene transfer

There was no significant difference in the frequency or amplitude of EPSCs or IPSCs in stratum radiatum interneurons following $\alpha 7$ gene transfer (data not shown). In stratum radiatum interneurons with $\alpha 7$ gene transfer ($n=8$) the frequency of EPSCs was 1.8 ± 0.9 Hz, and in control cells ($n=9$) it was 1.3 ± 0.4 Hz. The amplitude of spontaneous EPSCs in GFP+ stratum radiatum interneurons following $\alpha 7$ gene transfer was 15.0 ± 1.4 pA, and in control cells it was 13.2 ± 1.0 pA. The frequency of IPSCs in stratum radiatum interneurons with $\alpha 7$ gene transfer




Fig. 5 – Group data showing the effects of 4OH-GTS-21 (black bar), 5-HI + 4OH-GTS-21 (light gray bar), 5-HI + 4OH-GTS-21 + TTX (hatched bar), and 5-HI + 4OH-GTS-21 + MLA (white bar) on the frequency (A) and amplitude (B) of spontaneous IPSCs in hippocampal CA1 pyramidal cells from normal animals. * indicates statistically significant changes from baseline amplitude and/or frequency where $p < 0.05$ and ** where $p < 0.01$.

Table 2 – IPSCs

Treatment	Baseline Hz	Treatment Hz	t-test	Baseline pA	Treatment pA	t-test	N
Normals							
4OH-GTS-21	1.8±0.3	1.9±0.3	–	33.9±3.1	28±1.9	–	8
5-HI+4OH-GTS-21	2.2±0.2	3.2±0.4	**	24.5±2.5	45.2±7.9	**	13
5-HI+4OH-GTS-21+TTX	2.8±0.2	4.4±0.6	*	23.2±1.8	20.1±2.1	–	5
5-HI+4OH-GTS-21+MLA	1.5±0.8	2.2±0.8	–	38.6±6.1	50.2±12.4	–	3
Gene transfer							
4OH-GTS-21	2.1±0.2	2.2±0.2	–	30.9±2.8	23.6±2.0	–	8
5-HI+4OH-GTS-21	2.3±0.22	2.9±0.4	**	25.4±4.0	53.0±12.1	**	8
5-HI+4OH-GTS-21+MLA	2.5±0.4	3.2±0.4	–	31.9±6.7	59.3±13.1	*	4

Raw data showing the frequencies and amplitudes of IPSCs ± SEM in CA1 pyramidal cells from normal animals and animals that received $\alpha 7$ gene transfer in baseline and drug treatment conditions. * $p < 0.05$, ** $p < 0.01$.

($n=9$) was 12.4 ± 1.1 Hz, and in control cells ($n=9$) it was 13.4 ± 2.2 Hz. The amplitude of IPSCs was 24.7 ± 2.5 pA in control cells and 21.0 ± 2.7 pA in stratum radiatum interneurons with $\alpha 7$ gene transfer.

3. Discussion

In the present report, 5-HI and 5-HI+4OH-GTS-21 significantly increased the frequency of spontaneous APV- and NBQX-sensitive EPSCs. Also, 5-HI+4OH-GTS-21 significantly increased the frequency and amplitude of spontaneous bicuculline-sensitive IPSCs, although 4OH-GTS-21 produced no significant effect when applied alone. The 5-HI+4OH-GTS-21 induced increase in EPSC frequency was blocked by TTX, but 5-HI induced effects were not blocked by application of MLA. In contrast, the 5-HI+4OH-GTS-21 increase in IPSC frequency was not blocked or reduced by application of TTX or MLA, but 5-HI+4OH-GTS-21 mediated increases in IPSC amplitudes were blocked by TTX and reduced by MLA in normal animals. In cells from $\alpha 7$ gene transfer animals, however, MLA was not effective in reducing increases in IPSC amplitudes. Finally, $\alpha 7$ gene transfer did not alter the magnitude of effects produced by 5-HI or 5-HI+4OH-GTS-21, and did not affect the baseline frequency or amplitude of synaptic events in stratum radiatum interneurons. Thus, indirect measures of $\alpha 7$ nAChR function were not significantly enhanced by combining $\alpha 7$ gene transfer with either agonist activation or positive allosteric modulation of $\alpha 7$ nAChRs.

When applied alone, or combined with 4OH-GTS-21, 5-HI increased the frequency of EPSCs. Because there was no frequency or amplitude increase produced with the selective $\alpha 7$ agonist 4OH-GTS-21, and no significant difference between the magnitude of increases produced by 5-HI when applied alone and with 4OH-GTS-21 (5-HI+4OH-GTS-21), it appears that the effects of 5-HI+4OH-GTS-21 resulted from the actions of 5-HI alone. Since the 5-HI+4OH-GTS-21 mediated increase in EPSC frequency was blocked with TTX but unchanged by MLA, our data suggest that these effects were not mediated by direct activation of $\alpha 7$ nAChRs, but rather by increases in action potential-dependent transmitter release. This is consistent with studies that have shown 5-HI mediated increases in glutamate release, the amplitude of population spikes, and the amplitude of evoked excitatory postsynaptic potentials in the hippocampus (Mannaioni et al., 2003). However, to our

knowledge, this is the first report showing that 5-HI potentiates the frequency of spontaneous EPSCs in CA1 pyramidal cells.

Our results are also consistent with studies showing that 5-HI increases the frequency and amplitude of spontaneous GABAergic inhibitory postsynaptic currents (IPSCs) (Mannaioni et al., 2003; Mok and Kew, 2006). Because TTX did not reduce the 5-HI+4OH-GTS-21 mediated frequency increase in IPSCs (Fig. 5A), our results suggest that 5-HI+4OH-GTS-21 may have increased IPSC frequencies through mechanisms not related to action potential-dependent transmitter release, perhaps ones related to cell excitability, even though we measured no significant changes in holding currents measured at the cell body. Because TTX did not reduce the 5-HI+4OH-GTS-21 mediated frequency increase in IPSCs (Fig. 5A), increased IPSC frequencies in response to 5-HI+4OH-GTS-21 are not related to action potential-dependent transmitter release, and the absence of significant changes in holding currents measured at the cell body (data not shown) makes it unlikely that cell excitability was altered. Further, because TTX blocked, and MLA reduced the 5HI+4OH-GTS-21 mediated increase in IPSC amplitudes (Fig. 5B), this effect likely involves both activation of $\alpha 7$ nAChRs and increases in action potential-dependent GABA release. Curiously however, MLA did not significantly reduce the 5-HI+4OH-GTS-21 potentiation of IPSC amplitudes in cells from animals that received $\alpha 7$ gene transfer. Any number of functional changes or potential changes involving synaptic architecture resulting from gene transfer could have produced this difference.

With the exception of a reduction in the potentiation of IPSC amplitudes mentioned above, none of the effects of 5-HI or 5-HI+4OH-GTS-21 was reversed or completely blocked by MLA. In this regard, our findings are consistent with Mannaioni et al., 2003, who showed that 5-HI potentiation of IPSCs in CA1 pyramidal cells was not sensitive to preincubation with MLA. In contrast, Mok and Kew, 2006 showed that 5-HI potentiation of IPSCs in stratum radiatum interneurons was occluded by pretreatment with MLA, TTX, and nicotine (hypothesized to desensitize receptors). Taken together, our results and those of Mannaioni et al., 2003 and Mok and Kew, 2006 demonstrate that 5-HI facilitates both excitatory and inhibitory neurotransmission. Further, they show that while the 5-HI mediated potentiation of IPSCs in stratum radiatum interneurons can be mediated by activation of $\alpha 7$ nAChRs, IPSC potentiation in CA1 pyramidal cells is probably mediated




Fig. 6 – Group data showing the effects of 4OH-GTS-21 (black bar), 5-HI + 4OH-GTS-21 (light gray bar), and 5-HI + 4OH-GTS-21 + MLA (white bar) on the frequency (A) and amplitude (B) of spontaneous IPSCs in hippocampal pyramidal cells from animals that received $\alpha 7$ gene transfer. * indicates statistically significant changes from baseline amplitude and/or frequency where $p < 0.05$ and ** where $p < 0.01$.

by both an increase in action potential-dependent transmitter release and activation of $\alpha 7$ nAChRs.

Because the activation of $\alpha 7$ nAChRs has been shown to enhance transmitter release at various synapses (Gray et al., 1996; MacDermott et al., 1999; McGehee et al., 1995; Seguela et al., 1993), we tested the hypothesis that $\alpha 7$ gene transfer would change the frequency and/or amplitude of spontaneous IPSCs and EPSCs. Further, we tested the hypothesis that the magnitude of effects produced by 5-HI and 4OH-GTS-21 would be greater following $\alpha 7$ nAChR gene transfer. Although we found no change in the baseline frequency and amplitude of EPSCs and IPSCs in CA1 pyramidal cells and stratum radiatum interneurons following $\alpha 7$ nAChR gene transfer, the coupling

of $\alpha 7$ nAChRs at various synapses is well established. For example, it has been shown that nAChRs occur presynaptically and postsynaptically at GABAergic synapses (Fabian-Fine et al., 2001) and more recently that $\alpha 7$ nAChR activation inhibits evoked GABA_A receptor-mediated currents and increases the frequency of spontaneous miniature IPSCs in hippocampal interneurons (Wanaverbecq et al., 2007). Accordingly, it seems possible that changes in evoked IPSCs or miniature IPSCs could have occurred following $\alpha 7$ gene transfer but were not resolved in the current study. Further, because studies implementing ultrastructural resolution have shown that $\alpha 7$ nAChRs tend to occur away from sites of ACh release, ACh may provide a more diffuse type of hippocampal input (Descarries et al., 2004), making certain effects produced by $\alpha 7$ gene transfer undetectable within the present methodologies.

We hypothesized that a combination of an $\alpha 7$ -selective agonist, allosteric modulator, and $\alpha 7$ gene transfer would be ideal for inducing maximum activation of $\alpha 7$ nAChRs in the hippocampal slice. However, our data suggest that although combining these methods may have amplified certain physiological functions of $\alpha 7$ nAChRs, we were unable to elucidate a role for hippocampal $\alpha 7$ nAChRs in excitatory and inhibitory synaptic transmission with this combination of treatments. Moreover, no effects on synaptic transmission could be resolved with the $\alpha 7$ -selective agonist 4OH-GTS-21 at the concentration tested, although this was within a concentration range shown to be efficacious with in vitro measures of GTS-21 mediated cytoprotection (Ren et al., 2005). Furthermore, the $\alpha 7$ allosteric modulator 5-HI appeared to have affected synaptic transmission primarily via non- $\alpha 7$ -related mechanisms. Interestingly, consistent with this finding, more recent results from our laboratory show that 5-HI does not increase whole-cell currents produced by focal somatic application of 4OH-GTS-21 in stratum radiatum interneurons, although other positive modulators can increase 4OH-GTS-21 evoked responses and 5-HI can increase $\alpha 7$ responses to other partial agonists (Lopez et al., 2007 unpublished data). These findings suggest a complex interaction between partial agonists and positive modulators of $\alpha 7$ nAChRs that should be considered in future studies seeking experimental effects by combining these compounds. Finally, $\alpha 7$ gene transfer produced no significant effects on spontaneous synaptic events, although we previously showed that direct measures of $\alpha 7$ function (e.g. the magnitude of ACh-evoked currents) are in fact significantly affected by $\alpha 7$ gene transfer (Ren et al., 2007b). Thus, establishing a definitive role for the physiological function of $\alpha 7$ nAChRs in hippocampal synaptic transmission remains elusive but may be accomplished in future studies investigating other mechanisms or substrates.

4. Experimental procedures

4.1. rAAV8 preparation

The recombinant adeno-associated virus serotype 8 (rAAV8) hybrid vector was used for transduction of the hippocampus because of its ability to achieve gene transfer within large tissue volumes (Klein et al., 2006). The rAAV8 vector was prepared and quantified using the methods of Zolotukhin et al. (1999), Klein et al. (2006), and Ren et al. (2007a,b).

Expression of green fluorescent protein (GFP) or rat $\alpha 7$ nAChRs was driven by a chicken β -actin promoter containing the human cytomegalovirus enhancer. Plasmids were propagated in SURE cells (Stratagene, LaJolla, CA, USA) and CsCl-purified. Briefly, 70% confluent human embryonic kidney 293 cells were transfected by the calcium phosphate method with AAV terminal repeat-containing GFP or rat $\alpha 7$ nicotinic receptor plasmid in equal molar ratios with the rAAV8 helper plasmid. After 3 days, cells and media were harvested and centrifuged at 3000 $\times g$. The pellets were resuspended in a solution of 50 mM Tris, pH 8.3 and 150 mM NaCl, then freeze-thawed three times. The resulting suspension was put through a discontinuous iodixanol gradient followed by a Q-sepharose column (Sigma Chemicals, St. Louis, MO, USA) to purify rAAV8. Vector doses were expressed as genomic particles (gp).

4.2. Hippocampal injections of rAAV8-rat $\alpha 7$ and rAAV8-GFP

All procedures involving animals were approved by the University of Florida Institutional Animal Care and Use Committee and were in accord with the NIH Guide for the Care and Use of Laboratory Animals. Male Sprague Dawley rats were maintained on a 12 h day night light cycle and had food and water available ad libitum until used in experiments. At 18–22 days of age they were placed in an isoflurane containing chamber for 5 min prior to mounting on a stereotaxic frame. Lack of withdrawal reflex indicated readiness for making the initial incision, which followed the midline suture from just rostral to bregma, continuing back to lambda. A small hole (0.5 mm diameter) was drilled 3.6 mm caudal to bregma and 2.2 mm lateral to the midline. A 27-gauge needle attached to a 10 μ l Hamilton syringe was lowered into the hippocampus and stopped at a depth 2.6–3.0 mm ventral to the skull surface. Equal volumes of rAAV8-rat $\alpha 7$ and rAAV8-GFP vectors [2–3 μ l total volume (10^{10} gp)] were delivered at 0.2 μ l/min using a CMA/100 microinjection pump (CMA/Microdialysis, Solna, Sweden). The incisions were closed using surgical nylon and treated with betadine. Animals recovered from anesthesia on a heated blanket and were monitored for signs of pain. Additional analgesic injections were administered as necessary. Animals were kept for 7–10 days in standard housing prior to use for electrophysiological experiments.

4.3. Patch-clamping in hippocampal slices

Animals were anesthetized with halothane (Halocarbon Laboratories, River Edge, NJ) and swiftly decapitated. Transverse (300 μ m) whole brain slices were prepared using a vibratome (Pelco, Redding, CA) and a high Mg^{2+} /low Ca^{2+} ice-cold artificial cerebral spinal fluid (ACSF) containing (in mM) 124 NaCl, 2.5 KCl, 1.2 NaH_2PO_4 , 2.5 $MgSO_4$, 10 D-glucose, 1 $CaCl_2$, and 25.9 $NaHCO_3$ saturated with 95% O_2 –5% CO_2 . Prior to sectioning, a single-edge razor was used to make a longitudinal cut that separated the two hemispheres which allowed for keeping the injected and un-injected sides separate. Slices were incubated at 30 $^\circ C$ for 30 min and remained at room temperature until they were transferred to a submersion chamber (Warner Instruments, Hamden, CT) for

recording. Slices were perfused (2 ml/min) with normal ACSF containing (in mM) 126 NaCl, 3 KCl, 1.2 NaH_2PO_4 , 1.5 $MgSO_4$, 11 D-glucose, 2.4 $CaCl_2$, 25.9 $NaHCO_3$, and 0.004–0.008 atropine sulfate saturated with 95% O_2 –5% CO_2 at 30 $^\circ C$. Cells were visualized with infrared differential interference contrast (IR DIC) using a Nikon E600FN microscope that was equipped with fluorescence for viewing GFP in injected cells. In addition to confirmation of fluorescence, functional expression was confirmed in some cells using somatic application of ACh (1 mM) which in all cases produced evoked responses (data not shown). Whole cell patch-clamp recordings were made with glass pipettes (3–5 $M\Omega$) containing an internal solution of (in mM) 125 K-gluconate, 1 KCl, 0.1 $CaCl_2$, 2 $MgCl_2$, 1 EGTA, 2 Mg-ATP, 0.3 Na_3GTP , and 10 HEPES (pH 7.3 using KOH). To record inhibitory post synaptic currents (IPSCs), cells were held at 0 mV using an internal solution of (in mM): 140 CsMeSO₃, 8 NaCl, 1 $MgCl_2$, 0.2 EGTA, 10 HEPES, 2 Mg-ATP, 0.3 Na-GTP, and 5 QX-314 (pH 7.3 using CsOH). Prior to data collection, all cells were held at –70 mV, and –10 mV/10 ms test pulses were used to determine series resistance (R_a), input resistance (R_m) and whole-cell capacitance (C_m). Cells with series resistances >60 $M\Omega$ or those requiring holding currents >250 pA were not included in the final analyses.

4.4. Drug application and data analyses

Bath applications of all drugs were achieved by introducing them into the ACSF using a syringe pump (Kd Scientific, Holliston, MA) loaded with a concentrated stock solution diluted to the final concentration in the perfusion line prior to entering the recording chamber. Baseline recordings of spontaneous synaptic events were made for a period of 2 min. Each drug was washed in for 6 min, and the following 2 min of recording was used to measure drug effects. Drugs were applied sequentially or together as indicated in the text. Since the effects of 1 mM 5-HI combined with 5 μ M 4OH-GTS-21 (5-HI+4OH-GTS-21) were not significantly different if the drugs were applied sequentially or together, in some cases data from each type of application were pooled together. 4OH-GTS-21 was synthesized and provided by Taiho Pharmaceuticals (Tokushima, Japan) and 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione (NBQX) was purchased from Tocris Cookson (Ellisville, MO). All other chemicals were obtained from Sigma (St. Louis, MO).

Signals were digitized using an Axon Digidata 1322A and sampled at 20 kHz on a Dell computer using Clampex version 8 or 9. Data analysis was done with Clampfit version 8 or 9, Excel 2000, and GraphPad/Prism version 3.02. Spontaneous synaptic events were analyzed using Clampfit event detection (threshold search in manual mode). Statistical analyses used paired and unpaired one-tailed Student's *t*-tests where appropriate. Data are expressed as mean \pm SEM.

Acknowledgments

The authors thank Clare Stokes for the technical assistance. This work was supported by the Evelyn F. McKnight Brain Research Foundation, NIH grants GM57481 to RLP and P01AG10485 to R.L.P. and E.M.M., and M.A.K.

REFERENCES

- Alkondon, M., Pereira, E.F., Barbosa, C.T., Albuquerque, E.X., 1997. Neuronal nicotinic acetylcholine receptor activation modulates gamma-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J. Pharmacol. Exp. Ther.* 283, 1396–1411.
- Arendash, G.W., Sengstock, G.J., Sanberg, P.R., Kem, W.R., 1995. Improved learning and memory in aged rats with chronic administration of the nicotinic receptor agonist GTS-21. *Brain Res.* 674, 252–259.
- Briggs, C.A., Anderson, D.J., Brioni, J.D., Buccafusco, J.J., Buckley, M.J., Campbell, J.E., Decker, M.W., Donnelly-Roberts, D., Elliott, R.L., Gopalakrishnan, M., Holladay, M.W., Hui, Y.H., Jackson, W.J., Kim, D.J., Marsh, K.C., O'Neill, A., Prendergast, M.A., Ryther, K.B., Sullivan, J.P., Americ, S.P., 1997. Functional characterization of the novel neuronal nicotinic acetylcholine receptor ligand GTS-21 in vitro and in vivo. *Pharmacol. Biochem. Behav.* 57, 231–241.
- Court, J., Spurden, D., Lloyd, S., McKeith, I., Ballard, C., Cairns, N., Kerwin, R., Perry, R., Perry, E., 1999. Neuronal nicotinic receptors in dementia with Lewy bodies and schizophrenia: alpha-bungarotoxin and nicotine binding in the thalamus. *J. Neurochem.* 73, 1590–1597.
- Dani, J.A., 2000. Properties underlying the influence of nicotinic receptors on neuronal excitability and epilepsy. *Epilepsia* 41, 1063–1065.
- Descarries, L., Mechawar, N., Aznavour, N., Watkins, K.C., 2004. Structural determinants of the roles of acetylcholine in cerebral cortex. *Prog. Brain Res.* 145, 45–58.
- Fabian-Fine, R., Skehel, P., Errington, M.L., Davies, H.A., Sher, E., Stewart, M.G., Fine, A., 2001. Ultrastructural distribution of the alpha7 nicotinic acetylcholine receptor subunit in rat hippocampus. *J. Neurosci.* 21, 7993–8003.
- Frazier, C.J., Strowbridge, B.W., Papke, R.L., 2003. Nicotinic receptors on local circuit neurons in dentate gyrus: a potential role in regulation of granule cell excitability. *J. Neurophysiol.* 89, 3018–3028.
- Freedman, R., Coon, H., Myles-Worsley, M., Orr-Urtreger, A., Olincy, A., Davis, A., Polymeropoulos, M., Holik, J., Hopkins, J., Hoff, M., Rosenthal, J., Waldo, M.C., Reimherr, F., Wender, P., Yaw, J., Young, D.A., Breese, C.R., Adams, C., Patterson, D., Adler, L.E., Leonard, S., Byerley, W., 1997. Linkage of a neurophysiological deficit in schizophrenia to a chromosome 15 locus. *Proc. Natl. Acad. Sci. U. S. A.* 94, 587–592.
- Gray, R., Rajan, A.S., Radcliffe, K.A., Yakehiro, M., Dani, J.A., 1996. Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature* 383, 713–716.
- Gronlien, J.H., Hakerud, M., Ween, H., Thorin-Hagene, K., Briggs, C.A., Gopalakrishnan, M., Malysz, J., 2007. Distinct profiles of alpha7 nAChR positive allosteric modulation revealed by structurally diverse chemotypes. *Mol. Pharmacol.* 72, 715–724.
- Guan, Z.Z., Zhang, X., Ravid, R., Nordberg, A., 2000. Decreased protein levels of nicotinic receptor subunits in the hippocampus and temporal cortex of patients with Alzheimer's disease. *J. Neurochem.* 74, 237–243.
- Gurley, D.A., Harris, E.W., Li, C., Johnson, E.C., Lanthorn, T.H., 2000. 5-hydroxyindole potentiates the nicotinic acetylcholine receptor alpha 7 subtype. *Society for Neuroscience Annual Meeting*, vol. 716.15. Society for Neuroscience, New Orleans, Louisiana.
- Hurst, R.S., Hajos, M., Raggenbass, M., Wall, T.M., Higdon, N.R., Lawson, J.A., Rutherford-Root, K.L., Berkenpas, M.B., Hoffmann, W.E., Piotrowski, D.W., Groppi, V.E., Allaman, G., Ogier, R., Bertrand, S., Bertrand, D., Arneric, S.P., 2005. A novel positive allosteric modulator of the alpha7 neuronal nicotinic acetylcholine receptor: in vitro and in vivo characterization. *J. Neurosci.* 25, 4396–4405.
- Ji, D., Dani, J.A., 2000. Inhibition and disinhibition of pyramidal neurons by activation of nicotinic receptors on hippocampal interneurons. *J. Neurophysiol.* 83, 2682–2690.
- Klein, R.L., Dayton, R.D., Leidenheimer, N.J., Jansen, K., Golde, T.E., Zweig, R.M., 2006. Efficient neuronal gene transfer with AAV8 leads to neurotoxic levels of tau or green fluorescent proteins. *Mol. Ther.* 13, 517–527.
- Krause, R.M., Buisson, B., Bertrand, S., Corringer, P.J., Galzi, J.L., Changeux, J.P., Bertrand, D., 1998. Ivermectin: a positive allosteric effector of the alpha7 neuronal nicotinic acetylcholine receptor. *Mol. Pharmacol.* 53, 283–294.
- Li, X., Rainnie, D.G., McCarley, R.W., Greene, R.W., 1998. Presynaptic nicotinic receptors facilitate monoaminergic transmission. *J. Neurosci.* 18, 1904–1912.
- MacDermott, A.B., Role, L.W., Siegelbaum, S.A., 1999. Presynaptic ionotropic receptors and the control of transmitter release. *Annu. Rev. Neurosci.* 22, 443–485.
- Mannaioni, G., Carpenedo, R., Moroni, F., 2003. 5-hydroxyindole causes convulsions and increases transmitter release in the CA1 region of the rat hippocampus. *Br. J. Pharmacol.* 138, 245–253.
- McGehee, D.S., Heath, M.J., Gelber, S., Devay, P., Role, L.W., 1995. Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science* 269, 1692–1696.
- Mesulam, M., 2004. The cholinergic lesion of Alzheimer's disease: pivotal factor or side show? *Learn. Mem.* 11, 43–49.
- Meyer, E.M., King, M.A., Meyers, C., 1998a. Neuroprotective effects of 2,4-dimethoxybenzylidene anabaseine (DMXB) and tetrahydroaminoacridine (THA) in neocortices of nucleus basalis lesioned rats. *Brain Res.* 786, 252–254.
- Meyer, E.M., Kuryatov, A., Gerzanich, V., Lindstrom, J., Papke, R.L., 1998b. Analysis of 3-(4-hydroxy, 2-methoxybenzylidene)anabaseine selectivity and activity at human and rat alpha-7 nicotinic receptors. *J. Pharmacol. Exp. Ther.* 287, 918–925.
- Mok, M.H.S., Kew, J.N., 2006. Excitation of rat hippocampal interneurons via modulation of endogenous agonist activity at the alpha7 nicotinic ACh receptor. *J. Physiol.* 574, 699–710.
- Mufson, E.J., Ginsberg, S.D., Ikonovic, M.D., DeKosky, S.T., 2003. Human cholinergic basal forebrain: chemoanatomy and neurologic dysfunction. *J. Chem. Neuroanat.* 26, 233–242.
- Ren, K., Puig, V., Papke, R.L., Itoh, Y., Hughes, J.A., Meyer, E.M., 2005. Multiple calcium channels and kinases mediate alpha7 nicotinic receptor neuroprotection in PC12 cells. *J. Neurochem.* 94, 926–933.
- Ren, K., King, M.A., Liu, J., Siemann, J., Altman, M., Meyers, C., Hughes, J.A., Meyer, E.M., 2007a. The alpha7 nicotinic receptor agonist 4OH-GTS-21 protects axotomized septohippocampal cholinergic neurons in wild type but not amyloid-overexpressing transgenic mice. *Neuroscience* 148, 230–237.
- Ren, K., Thinschmidt, J., Liu, J., Ai, L., Papke, R.L., King, M.A., Hughes, J.A., Meyer, E.M., 2007b. alpha7 Nicotinic receptor gene delivery into mouse hippocampal neurons leads to functional receptor expression, improved spatial memory-related performance, and tau hyperphosphorylation. *Neuroscience* 145, 314–322.
- Santos, M.D., Alkondon, M., Pereira, E.F., Aracava, Y., Eisenberg, H.M., Maelicke, A., Albuquerque, E.X., 2002. The nicotinic allosteric potentiating ligand galantamine facilitates synaptic transmission in the mammalian central nervous system. *Mol. Pharmacol.* 61, 1222–1234.
- Schildstrom, B., Svensson, H.M., Svensson, T.H., Nomikos, G.G., 1998. Nicotine and food induced dopamine release in the nucleus accumbens of the rat: putative role of alpha7 nicotinic receptors in the ventral tegmental area. *Neuroscience* 85, 1005–1009.
- Seguela, P., Wadiche, J., Dineley-Miller, K., Dani, J.A., Patrick, J.W., 1993. Molecular cloning, functional properties, and distribution

- of rat brain alpha 7: a nicotinic cation channel highly permeable to calcium. *J. Neurosci.* 13, 596–604.
- Teaktong, T., Graham, A., Court, J., Perry, R., Jaros, E., Johnson, M., Hall, R., Perry, E., 2003. Alzheimer's disease is associated with a selective increase in alpha7 nicotinic acetylcholine receptor immunoreactivity in astrocytes. *Glia* 41, 207–211.
- Thinschmidt, J.S., Frazier, C.J., King, M.A., Meyer, E.M., Papke, R.L., 2005. Septal innervation regulates the function of alpha7 nicotinic receptors in CA1 hippocampal interneurons. *Exp. Neurol.* 195, 342–352.
- Uteshev, V.V., Meyer, E.M., Papke, R.L., 2003. Regulation of neuronal function by choline and 4OH-GTS-21 through alpha 7 nicotinic receptors. *J. Neurophysiol.* 89, 1797–1806.
- Wanaverbecq, N., Semyanov, A., Pavlov, I., Walker, M.C., Kullmann, D.M., 2007. Cholinergic axons modulate GABAergic signaling among hippocampal interneurons via postsynaptic alpha 7 nicotinic receptors. *J. Neurosci.* 27, 5683–5693.
- Woodruff-Pak, D.S., Li, Y.T., Kem, W.R., 1994. A nicotinic agonist (GTS-21), eyeblink classical conditioning, and nicotinic receptor binding in rabbit brain. *Brain Res.* 645, 309–317.
- Zolotukhin, S., Byrne, B.J., Mason, E., Zolotukhin, I., Potter, M., Chesnut, K., Summerford, C., Samulski, R.J., Muzyczka, N., 1999. Recombinant adeno-associated virus purification using novel methods improves infectious titer and yield. *Gene Ther.* 6, 973–985.
- Zwart, R., De Filippi, G., Broad, L.M., McPhie, G.I., Pearson, K.H., Baldwinson, T., Sher, E., 2002. 5-hydroxyindole potentiates human alpha 7 nicotinic receptor mediated responses and enhances acetylcholine-induced glutamate release in cerebellar slices. *Neuropharmacology* 43, 374–384.